arXiv:1612.06660v3 [hep-th] 26 Feb 2017

KOBE-TH-16-09

Polyakov Loop in Non-covariant Operator Formalism

Makoto Sakamoto[(a) 1] and Kazunori Takenaga[(b) 2]

[(a)] *Department of Physics, Kobe University, Rokkodai Nada, Kobe, 657-8501 Japan*

[(b)] *Faculty of Health Science, Kumamoto Health Science University, Izumi-machi, Kita-ku, Kumamoto 861-5598, Japan*

Abstract

We discuss a Polyakov loop in non-covariant operator formalism which consists of only physical degrees of freedom at finite temperature. It is pointed out that although the Polyakov loop is expressed by a Euclidean time component of gauge fields in a covariant path integral formalism, there is no direct counterpart of the Polyakov loop operator in the operator formalism because the Euclidean time component of gauge fields is not a physical degree of freedom. We show that by starting with an operator which is constructed in terms of only physical operators in the non-covariant operator formalism, the vacuum expectation value of the operator calculated by trace formula can be rewritten into a familiar form of an expectation value of Polyakov loop in a covariant path integral formalism at finite temperature for the cases of axial and Coulomb gauge.

[1]E-mail: dragon@kobe-u.ac.jp
[2]E-mail: takenaga@kumamoto-hsu.ac.jp

1 Introduction

Gauge invariance is undoubtedly one of fundamental principles in particle physics. Non-local gauge invariant operators as well as local ones play an important role in gauge theory. An example of such non-local operators is a Wilson loop, which will be given by a line integral along a rectangular contour that one side is taken to be in a space-like direction and another side to be in a time direction. A non-trivial expectation value of the Wilson loop operator will provide a signal of "quark" confinement in non-abelian gauge theories [1].

Another example of non-local gauge invariant operators is a Polyakov loop, which is similar to a Wilson loop but is given by a line integral along a Euclidean time axis. In this paper, we will focus on the Polyakov loop operator whose explicit form is given by $\mathrm{tr}\,\mathcal{P}\exp\left(ig\int_0^\beta d\tau A_\tau\right)$, where $\mathcal{P}$ denotes a path-ordered symbol, β is an inverse temperature and A_τ is a Euclidean time component of gauge fields. The Polyakov loop is known as an order parameter of the confinement-deconfinement phase transition at finite temperature [2, 3]. Furthermore, it could provide an order parameter of gauge symmetry breakings [2, 4, 5, 6, 7].

In order to confirm that the Polyakov loop is a physical observable at finite temperature, one might verify that a zero mode of the Euclidean time component A_τ in the Polyakov loop cannot be eliminated by gauge transformations due to the periodicity with respect to the Euclidean time at finite temperature. However, the statement that the Polyakov loop is physical seems to be less obvious than we thought.[3]

A reason why the above statement is not so obvious is that the Minkowski time component A_0 of gauge fields[4] is not a dynamical degree of freedom in gauge theory. The A_0 component can be removed in an operator formalism, and then the A_0 degree of freedom disappears completely from the Hilbert space of the theory, as explained below in some detail. One might stress that the A_0 (or A_τ) component is included in a covariant path integral representation of gauge theory. However, the A_0 (or A_τ) degree of freedom turns out to be introduced as *an auxiliary field* in a path integral formalism (see Section 3).

Quantization of gauge theory cannot be performed in a straightforward fashion because it contains redundant gauge degrees of freedom due to gauge invariance. Several ways to quantize it have been known and are expected to be physically equivalent.[5]

One of reliable methods to quantize gauge theory is to remove all unphysical degrees of freedom by explicitly imposing a gauge fixing condition [10]. If the theory includes only

[3] One might notice that Wilson lines along non-contractible loops in extra dimensions [8] resemble a Polyakov loop along the Euclidean time axis. There seems formally no difference between them in a Euclidean path integral formulation. However, there is an important difference. A Polyakov loop becomes meaningless at zero temperature but Wilson lines do not. This is because the Euclidean time component of gauge fields can be removed completely by gauge transformations at zero temperature but the extra dimensional components cannot be.

[4] In this paper, we will use A_τ and A_0 for the Euclidean and Minkowski time components of gauge fields, respectively.

[5] We will not discuss the Gribov problem [9] in this paper. The problem is expected not to be related directly to our issue.

physical degrees of freedom, the quantization of the theory is rather straightforward, though the form of the Hamiltonian would be messy.

For instance, in the axial gauge $A_3 = 0$ [11], which will be discussed in this paper, the theory can be described only by the physical degrees of freedom A_1 and A_2 for the gauge fields because A_3 is taken to be zero by the axial gauge condition and A_0 is also removed by use of the equation of motion for A_0, which should be regarded as a constraint equation since it includes no time derivative of A_0.[6] Therefore, the gauge part of the Hilbert space in the axial gauge formalism can be spanned by the states $\{ \mid A_1, A_2 \rangle \}$, and any physical operators have to be constructed in terms of A_k $(k = 1, 2)$ and their conjugate momenta Π_k (and also fermion/scalar fields).

Now we can raise an issue of how to construct a Polyakov loop operator in the axial gauge operator formalism. Since the time component is completely eliminated from the spectrum in the operator formalism, it seems non-trivial to construct a Polyakov loop operator without A_0 or A_τ.

One of our main purposes is to present an explicit operator form corresponding to a Polyakov loop operator, in the operator formalism which contains only physical degrees of freedom: A_k, Π_k $(k = 1, 2)$ and matter fields. Another purpose of this paper is to show that an expectation value of the above operator given in the non-covariant operator formalism can be rewritten into a familiar form of an expectation value of a Polyakov loop in a covariant path integral formalism at finite temperature. In the rewriting, we will see where the Euclidean time component A_τ comes from and also clarify how the operator given by physical operators turns to the standard form of the Polyakov loop given by A_τ.

This paper is organized as follows. In the next section, we take the axial gauge to quantize a non-abelian gauge theory, and remove all the unphysical degrees of freedom from a Lagrangian. Then, we derive a Hamiltonian consisting of only physical ones. In Section 3, we first construct a finite temperature path integral representation in the axial gauge by use of the trace formula [13], and then rewrite it into a covariant path integral form with all the gauge degrees of freedom. We will see that the Euclidean time component of the gauge fields is recovered as a Gaussian auxiliary field. Our main results will be given in Section 4. We propose an operator that is expected to represent a Polyakov loop operator in the axial gauge and is written in terms of physical operators only. We show that the operator turns to the standard form of the Polyakov loop in a covariant Euclidean path integral formula at finite temperature. In Section 5, we reformulate the analysis in the Coulomb gauge. Section 6 is devoted to conclusions. Some details of calculations will be given in Appendices.

2 Hamiltonian in Axial Gauge

In this section, we present a canonical formulation of non-abelian gauge theory coupled with fermion in axial gauge. Although the results given in Sections 2 and 3 will not be new [10],

[6] If we take the temporal gauge $A_0 = 0$ [12], A_0 can be eliminated rather directly.

we shall explain them to make our discussions in Section 4 comprehensible. We consider an $SU(N)$ gauge theory whose Lagrangian is given by

$$\mathcal{L} = \frac{-1}{4} F^a_{\mu\nu} F^{a\mu\nu} + \bar{\psi} i \gamma^\mu D_\mu \psi, \tag{2.1}$$

where

$$F^a_{\mu\nu} \equiv \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + g f^{abc} A^b_\mu A^c_\nu, \quad D_\mu \psi \equiv \left(\partial_\mu - i g A^a_\mu T^a\right) \psi. \tag{2.2}$$

The representation of the fermion ψ under the $SU(N)$ gauge group can be arbitrary, but for simplicity, we take it to be the fundamental representation. Throughout this paper, we use the convention that repeated indices are summed over (otherwise stated) and $\mu, \nu, \cdots = 0, 1, 2, 3$ (or $\tau, 1, 2, 3$), $i, j = 1, 2, 3$, $k, l, \cdots = 1, 2$, $a, b, \cdots = 1, 2, \cdots, N^2 - 1$. The equation of motion for the gauge field A^a_μ followed from the Lagrangian (2.1) is

$$(D_\mu)^{ab} F^{b\mu\nu} + J_F^{a\nu} = 0, \tag{2.3}$$

where $J_F^{a\nu} \equiv g \bar{\psi} \gamma^\nu T^a \psi$ and $(D_\mu)^{ab} = \delta^{ab} \partial_\mu + g f^{acb} A^c_\mu$. We impose the axial gauge

$$A^a_3 = 0. \tag{2.4}$$

Then, we can formally solve the $\nu = 0$ component of the equation of motion for A^a_0 and obtain that

$$A^a_0 = (\mathcal{D}^{-1})^{ab} \Big((D_k \dot{A}_k)^b + J_F^{b0} \Big) \qquad (k = 1, 2). \tag{2.5}$$

Here the operator $(\mathcal{D}^{-1})^{ab}$ is formally defined as the inverse of $(\mathcal{D})^{ab} \equiv (D_k^2)^{ab} + \delta^{ab} (\partial_3)^2$.

We define the canonical momentum for the gauge field $A^a_k (k = 1, 2)$ by

$$\Pi_A^{ak} \equiv \frac{\partial \mathcal{L}(A_k, \psi)_{\rm axial}}{\partial \dot{A}^a_k}, \tag{2.6}$$

where $\mathcal{L}(A_k, \psi)_{\rm axial}$ is obtained by imposing the axial gauge on the Lagrangian (2.1) and eliminating A^a_0 in the Lagrangian (2.1) by using the equation (2.5), so that it is written in terms of only physical degrees of freedom. The explicit form of $\mathcal{L}(A_k, \psi)_{\rm axial}$ is given by Eq.(A.1) in the Appendix A, from which the canonical momentum for A^a_k is calculated as

$$\Pi_A^{ak} = (M)^{ab}_{kl} \dot{A}^b_l - (D_k \mathcal{D}^{-1} J_F^0)^a \qquad (l = 1, 2), \tag{2.7}$$

where

$$(M)^{ab}_{kl} = \delta^{ab} \delta_{kl} - (D_k \mathcal{D}^{-1} D_l)^{ab}. \tag{2.8}$$

By solving the equation (2.7) for $\dot{A}^a_k$, we have

$$\dot{A}^a_k = (M^{-1})^{ab}_{kl} \Big(\Pi_A^{bl} + (D_l \mathcal{D}^{-1} J_F^0)^b \Big), \tag{2.9}$$

where $(M^{-1})^{ab}_{kl}$ is defined as the inverse of $(M)^{ab}_{kl}$, whose explicit form is

$$(M^{-1})^{ab}_{kl} = \delta^{ab} \delta_{kl} + (D_k (\partial_3)^{-2} D_l)^{ab}. \tag{2.10}$$

It is straightforward to check $(MM^{-1})^{ab}_{km} = \delta^{ab}\delta_{km}$ by using $(\mathcal{D})^{ab} = (D_k^2)^{ab} + \delta^{ab}(\partial_3)^2$. As for the fermion ψ, we take the right derivative with respect to the Grassmann variable, and the canonical momentum for ψ is

$$\Pi_\psi \equiv \frac{\partial \mathcal{L}(A_k, \psi)_{\rm axial}}{\partial \dot{\psi}} = \bar{\psi} i\gamma^0. \tag{2.11}$$

Then, the Hamiltonian is

$$\begin{aligned}
\mathcal{H}_{\rm axial} &\equiv \Pi_A^{ak}\dot{A}_k^a + \Pi_\psi \dot{\psi} - \mathcal{L}(A_k,\psi)_{\rm axial} \\
&= \frac{1}{2}\Pi_A^{ak}(M_{kl}^{-1}\Pi_A^l)^a + \Pi_A^{ak}(M_{kl}^{-1}D_l\mathcal{D}^{-1}J_F^0)^a \\
&\quad + \frac{1}{2}(D_k\mathcal{D}^{-1}J_F^0)^a(M_{kl}^{-1}D_l\mathcal{D}^{-1}J_F^0)^a - \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a \\
&\quad + \frac{1}{4}F_{kl}^aF_{kl}^a + \frac{1}{2}\partial_3 A_k^a \partial_3 A_k^a - \bar{\psi} i\gamma^k D_k\psi - \bar{\psi} i\gamma^3\partial_3\psi.
\end{aligned} \tag{2.12}$$

In deriving the Hamiltonian, we have performed the partial integration with resect to $(M^{-1})^{ab}_{kl}$. The Hamiltonian is written in terms of only physical degrees of freedom A_k^a, Π_A^{ak}, ψ and Π_ψ as it should, though it is messy, including the non-local terms. It may be appropriate to mention here that one is able to rewrite (2.12) into another form presented by Eq.(A.6) in the Appendix A, in which the self-energy of the color charge densities of fermion manifestly appears. We shall use the Hamiltonian (2.12) in the discussions given below.

3 Trace Formula in Axial Gauge at Finite Temperature

Let us first define the vacuum expectation value of an operator $\hat{\mathcal{O}}$ at finite temperature by the trace formula,

$$\langle\hat{\mathcal{O}}\rangle_\beta \equiv {\rm Tr}\left(\hat{\mathcal{O}}\ {\rm e}^{-\beta\hat{H}}\right) \quad \text{with} \quad \hat{H} \equiv \int d^3\mathbf{x}\ \hat{\mathcal{H}}, \tag{3.1}$$

where β stands for the inverse of the temperature T. If one chooses a non-covariant gauge such as the axial or Coulomb gauge, the operator $\hat{\mathcal{O}}$ and the Hamiltonian $\hat{H}$ are written in terms of only physical degrees of freedom, *i.e.* $\hat{A}_k^a, \hat{\psi}$ and their canonical momenta, $\hat{\Pi}_A^{ak}, \hat{\Pi}_\psi$, so that the trace in Eq.(3.1) must be taken over the physical states alone, denoted here by $A_{\rm phys} = \{A_k^a, \psi\}$,

$$\begin{aligned}
&{\rm Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})\ {\rm e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})}\right)_{\rm gauge} \\
&= \int \mathcal{D}A_{\rm phys}\langle A_{\rm phys}|\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, , \hat{\Pi}_\psi, \hat{\psi})\ {\rm e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})}|A_{\rm phys}\rangle_{\rm gauge}.
\end{aligned} \tag{3.2}$$

It should be understood that bosons (fermions) must obey the (anti) periodic boundary condition for the Euclidean time direction because of the quantum statistics at finite temperature.

Our aim is to find the operator $\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})$ which satisfies

$$\mathrm{Tr}\left(\mathcal{O}(\hat{A}_k^a, \hat{\psi}, \hat{\Pi}_A^{ak}, \hat{\Pi}_\psi)\mathrm{e}^{-\beta H(\hat{A}_k^a, \hat{\psi}, \hat{\Pi}_A^{ak}, \hat{\Pi}_\psi)}\right)_{\mathrm{gauge}} = \int \mathcal{D}A_\mu \mathcal{D}\bar{\psi}\mathcal{D}\psi \ \det(M_{\mathrm{FP}}^{\mathrm{gauge}}) \prod_{x,a} \delta(\chi_{\mathrm{gauge}}^a(x))$$

$$\times \ \mathrm{tr}\Big(\mathcal{P} \ \mathrm{e}^{ig\int_0^\beta d\tau A_\tau(\tau, \mathbf{x}_0)}\Big) \ \exp\int_0^\beta d\tau \int d^3\mathbf{x}\Big\{\frac{-1}{4}F_{\mu\nu}^a F_{\mu\nu}^a - \bar{\psi}i\gamma_\mu D_\mu \psi\Big\}, \tag{3.3}$$

where $M_{\mathrm{FP}}^{\mathrm{gauge}}$ and $\delta(\chi_{\mathrm{gauge}}^a(x))$ are the Faddeev-Popov determinant and the gauge condition for the chosen gauge, respectively and $\mu, \nu = \tau, 1, 2, 3$. We shall consider the cases of the axial and Coulomb gauge in this paper. The right-hand-side of Eq.(3.3) is nothing but the vacuum expectation value of the Polyakov loop operator in the covariant path integral form under the chosen gauge condition. It should be noticed that the left-hand-side of Eq.(3.3) is given by the trace formula in the non-covariant operator formalism by use of the physical degrees of freedom alone, not including A_τ, especially that the operator $\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})$ has to be constructed by the physical operators only. Therefore, it seems to be a non-trivial problem whether or not there exists such an operator $\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})$ satisfying the equation (3.3). Furthermore, we need to clarify where the unphysical degree of freedom A_τ comes from in deriving the right-hand-side of Eq.(3.3) from the left-hand-side of it, and manage the path ordered product of the Polyakov loop in the trace formula.

Before we construct the operator $\hat{\mathcal{O}}$ explained above, let us first study the partition function at finite temperature, say, $\hat{\mathcal{O}} = \hat{\mathbf{1}}$ in the case of the axial gauge. In this case what we would like to show is that for the Hamiltonian (2.12) the equation,

$$\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})}\right)_{\mathrm{axial}}$$
$$= \int \mathcal{D}A_\mu \mathcal{D}\bar{\psi}\mathcal{D}\psi \ \det(\partial_3) \prod_{x,a} \delta(A_3^a) \ \exp\int_0^\beta d\tau \int d^3\mathbf{x}\Big\{\frac{-1}{4}F_{\mu\nu}^a F_{\mu\nu}^a - \bar{\psi}i\gamma_\mu D_\mu \psi\Big\} \tag{3.4}$$

holds where the right-hand-side of Eq.(3.4) is the standard path integral representation for the partition function at finite temperature in the axial gauge. This may be too pedagogical, but helpful for the later discussions. Using the completeness relation for the canonical momenta Π_A^{ak} and Π_ψ, the left-hand-side of Eq.(3.4) can be written in the path integral form as

$$\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})}\right)_{\mathrm{axial}}$$
$$= \int \mathcal{D}\Pi_A^{ak}\mathcal{D}A_k^a\mathcal{D}\Pi_\psi\mathcal{D}\psi \ \exp\int_0^\beta d\tau \int d^3\mathbf{x}\Big\{i\Pi_A^{ak}\dot{A}_k^a + i\Pi_\psi\dot{\psi} - \mathcal{H}_{\mathrm{axial}}\Big\}, \tag{3.5}$$

where $\mathcal{H}_{\mathrm{axial}}$ is given by Eq. (2.12) and $\dot{A}_k^a, \dot{\psi}$ stands for the derivative with respect to the Euclidean time τ, $\dot{A}_k^a = \partial_\tau A_k^a, \dot{\psi} = \partial_\tau \psi$. This notation will be used hereafter. Let us note that the imaginary unit " i " in front of $\Pi_A^{ak}\dot{A}_k^a$ and $\Pi_\psi\dot{\psi}$ does not come from the Euclideanization, but from taking the inner product of the "coordinate" $\phi(\mathbf{x})$ and the "momentum" $\Pi_\phi(\mathbf{x})$,

$$\langle\phi|\Pi_\phi\rangle \propto \exp\Big\{i\int d^3\mathbf{x} \ \Pi_\phi(\mathbf{x})\phi(\mathbf{x})\Big\}. \tag{3.6}$$

The exponent of the right-hand-side of Eq.(3.5) is quadratic with respect to Π_A^{ak}, whose explicit form is given by Eq.(A.7) in the Appendix A, so that we can perform the Gaussian integration $\mathcal{D}\Pi_A^{ak}$. Then, we arrive at

$$\begin{aligned}
&\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{axial}} = \int \mathcal{D}A_k^a\mathcal{D}\Pi_\psi\mathcal{D}\psi\ \det{}^{-\frac{1}{2}}(M^{-1})\ \exp\ \int_0^\beta d\tau \int d^3\mathbf{x} \\
&\times\Big\{-\frac{1}{2}\dot{A}_k^a\dot{A}_k^a - \frac{1}{2}(D_k\dot{A}_k)^a(\mathcal{D}^{-1}D_l\dot{A}_l)^a - i\dot{A}_k^a(D_k\mathcal{D}^{-1}J_F^0)^a + \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a \\
&\quad - \frac{1}{4}F_{kl}^aF_{kl}^a - \frac{1}{2}\partial_3A_k^a\partial_3A_k^a + i\Pi_\psi\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3\partial_3\psi\Big\}. \qquad (3.7)
\end{aligned}$$

In order to restore the A_τ degree of freedom, let us consider the Gaussian integral given by

$$\begin{aligned}
1 = &\int \mathcal{D}A_\tau^a\ \det{}^{\frac{1}{2}}(\mathcal{D})\ \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{-\frac{1}{2}\Big(iA_\tau^a - (\mathcal{D}^{-1})^{ac}\big(i(D_k\dot{A}_k)^c + J_F^{c0}\big)\Big) \\
&\times(\mathcal{D})^{ad}\Big(iA_\tau^d - (\mathcal{D}^{-1})^{de}\big(i(D_l\dot{A}_l)^e + J_F^{e0}\big)\Big)\Big\}. \qquad (3.8)
\end{aligned}$$

By inserting the equation (3.8) into Eq.(3.7), some of the terms in the exponent in Eqs.(3.7) and (3.8) are cancelled each other, as shown in the Appendix A.2, where we explicitly present the expansion of the exponent in the Gaussian integral (3.8) by Eq.(A.9). Then, we obtain that

$$\begin{aligned}
&\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{axial}} = \int \mathcal{D}A_\tau^a\mathcal{D}A_k^a\mathcal{D}\Pi_\psi\mathcal{D}\psi\ \det{}^{-\frac{1}{2}}(M^{-1})\det{}^{\frac{1}{2}}(\mathcal{D}) \\
&\times\ \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{-\frac{1}{2}\dot{A}_k^a\dot{A}_k^a - \frac{1}{4}F_{kl}^aF_{kl}^a - \frac{1}{2}\partial_3A_k^a\partial_3A_k^a \\
&-\frac{1}{2}(D_kA_\tau)^a(D_kA_\tau)^a - \frac{1}{2}\partial_3A_\tau^a\partial_3A_\tau^a + \dot{A}_k^a(D_kA_\tau)^a \\
&+i\Pi_\psi\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3\partial_3\psi + iA_\tau^aJ_F^{a0}\Big\}. \qquad (3.9)
\end{aligned}$$

Let us note that A_τ^a is recovered as the auxiliary field through the Gaussian integral (3.8). We finally introduce the factor,

$$1 = \int \mathcal{D}A_3^a\prod_{x,a}\delta(A_3^a) \qquad (3.10)$$

in order to restore the A_3^a degree of freedom. By inserting it into Eq.(3.9), the exponent in Eq.(3.9) is summarized into the covariant form of the Lagrangian thanks to the delta function in Eq.(3.10). And the determinants in Eq.(3.9) are evaluated as

$$\det{}^{-\frac{1}{2}}(M^{-1})\det{}^{\frac{1}{2}}(\mathcal{D}) = \det(\partial_3), \qquad (3.11)$$

which is shown in the Appendix C.1. Hence, we finally obtain that

$$\begin{aligned}
&\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{axial}} \\
= &\int \mathcal{D}A_\mu^a\mathcal{D}\bar{\psi}\mathcal{D}\psi\ \det(\partial_3)\prod_{x,a}\delta(A_3^a)\ \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{\frac{-1}{4}F_{\mu\nu}^aF_{\mu\nu}^a - \bar{\psi}i\gamma_\mu D_\mu\psi\Big\}. \qquad (3.12)
\end{aligned}$$

where we have used $\mathcal{D}\Pi_\psi = \mathcal{D}\bar{\psi}$ and defined $\gamma_\tau \equiv -i\gamma^0$. This is the equation (3.4) that we would like to prove.

4 Polyakov Loop in Axial Gauge

Let us discuss the Polyakov loop operator in the axial gauge which is one of our main purposes in the paper. As explained in the introduction, the gauge part of the Hilbert space in the axial gauge does not contain the gauge field A_τ, from which the Polyakov loop operator is usually defined. Thus, the problem is how to construct the operator written in terms of only physical degrees of freedom in the axial gauge, corresponding to the Polyakov loop operator. Our aim is to present an explicit form of the operator $\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})$ which satisfies

$$\begin{aligned} & \mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})}\right)_{\mathrm{axial}} \\ = & \int \mathcal{D}A_\mu^a \mathcal{D}\bar{\psi}\mathcal{D}\psi\ \det(\partial_3) \prod_{x,a} \delta(A_3^a)\ \mathrm{tr}\Big(\mathcal{P}\ \exp\Big\{ig\int_0^\beta d\tau A_\tau(\tau, \mathbf{x}_0)\Big\}\Big) \\ & \times \exp\int_0^\beta d\tau \int d^3\mathbf{x}\Big\{\frac{-1}{4}F_{\mu\nu}^a F_{\mu\nu}^a - \bar{\psi}i\gamma_\mu D_\mu \psi\Big\}, \end{aligned} \tag{4.1}$$

where the operator $\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})$ must be written in terms of only physical degrees of freedom in the axial gauge. The trace of the left-hand-side is taken over the physical state alone in the axial gauge where there is no A_τ^a degree of freedom. The right-hand-side is, of course, the vacuum expectation value of the Polyakov loop operator in the covariant path integral form with the axial gauge.

In our trials to find operators satisfying the equation (4.1), we have observed that the non-abelian nature of the path-ordered product of the Polyakov loop in Eq.(4.1) is an obstacle to rewrite the expectation value of the operator $\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})$ in the trace formula of the left-hand-side of Eq.(4.1) into the covariant path integral form in the right-hand-side of Eq.(4.1). Thus, it seems to be necessary to transform the right-hand-side of Eq.(4.1) into a path integral form without path-ordered product.

To this end, let us consider the gauge transformations such as

$$A_\tau^a(\tau, \mathbf{x})T^a = U(\tau)\Big(A'^a_\tau(\tau, \mathbf{x})T^a + \frac{i}{g}\partial_\tau\Big)U^\dagger(\tau), \tag{4.2}$$

$$A_i^a(\tau, \mathbf{x})T^a = U(\tau)A'^a_i(\tau, \mathbf{x})T^a U^\dagger(\tau), \tag{4.3}$$

$$\psi(\tau, \mathbf{x}) = U(\tau)\psi'(\tau, \mathbf{x}), \tag{4.4}$$

where the unitary matrix $U(\tau)$ is assumed to depend only on the Euclidean time τ with the periodic boundary condition $U(\tau + \beta) = U(\tau)$ in order not to contradict with the quantum statistics at finite temperature. We notice that there is no inhomogeneous term in the right-hand-side of Eq.(4.3) because $U(\tau)$ does not depend on the spatial coordinate $\mathbf{x}$. If we formally

write the delta functions $\prod_{x,a}\delta(A_3^a(\tau,\mathbf{x}))$ as $\prod_x\delta(A_3(\tau,\mathbf{x}))$ with $A_3 = A_3^a T^a$, it will be obvious that the delta functions are invariant under the gauge transformation (4.3), *i.e.*

$$\prod_x \delta(A_3(\tau,\mathbf{x})) = \prod_x \delta(A_3'(\tau,\mathbf{x})). \tag{4.5}$$

We choose the unitary matrix $U(\tau)$ to diagonalize the Polyakov loop by the gauge transformation (4.2) as follows: The path-ordered product $\mathcal{P}\ \mathrm{e}^{ig\int_0^\beta d\tau A_\tau^a(\tau,\mathbf{x}_0)T^a}$ transforms under the gauge transformation (4.2) as

$$\begin{aligned}\mathcal{P}\ \mathrm{e}^{ig\int_0^\beta d\tau A_\tau^a(\tau,\mathbf{x}_0)T^a} &= U(\beta)\left[\mathcal{P}\ \mathrm{e}^{ig\int_0^\beta d\tau A'^a_\tau(\tau,\mathbf{x}_0)T^a}\right]U^\dagger(0)\\ &= U(0)\left[\mathcal{P}\ \mathrm{e}^{ig\int_0^\beta d\tau A'^a_\tau(\tau,\mathbf{x}_0)T^a}\right]U^\dagger(0).\end{aligned} \tag{4.6}$$

Since $\mathcal{P}\ \mathrm{e}^{ig\int_0^\beta d\tau A'^a_\tau(\tau,\mathbf{x}_0)T^a}$ is a unitary matrix fixed at $\mathbf{x}_0$, it can be diagonalized by a unitary matrix $U(0)$. Then, we can rewrite the Polyakov loop into the form

$$\begin{aligned}\mathrm{tr}\Big(\mathcal{P}\ \mathrm{e}^{ig\int_0^\beta d\tau A_\tau^a(\tau,\mathbf{x}_0)T^a}\Big) &= \mathrm{tr}\Big(\exp\Big\{g\int_0^\beta d\tau A'^{\tilde{a}}_\tau(\tau,\mathbf{x}_0)T^{\tilde{a}}\Big\}\Big)\\ &= \sum_{\alpha=1}^D \exp\Big\{ig\int_0^\beta d\tau A'^{\tilde{a}}_\tau(\tau,\mathbf{x}_0)(T^{\tilde{a}})_{\alpha\alpha}\Big\},\end{aligned} \tag{4.7}$$

where $\{T^{\tilde{a}}, \tilde{a}=1,2,\cdots,N-1\}$ are generators of the Cartan subalgebra of $SU(N)$ with the diagonal form

$$(T^{\tilde{a}})_{\alpha\beta} = (T^{\tilde{a}})_{\alpha\alpha}\delta_{\alpha\beta} \qquad (\alpha,\beta = 1,2,\cdots,D). \tag{4.8}$$

Here, D stands for the dimension of the representation of $\{T^a\}$. ($D=N$ for the fundamental representation.) Thus, the Polyakov loop turns out to reduce to the one defined for the $U(1)$ subgroups in the $SU(N)$ gauge group. Thanks to the reduction, the trace in the original expression (4.1) is replaced by the summation with respect to the abelian part of the Polyakov loop in Eq.(4.7) (note that $(T^{\tilde{a}})_{\alpha\alpha}$ is just a number) and one does not need to take care of the path-ordered integral because of the abelian nature of the $U(1)$ gauge group.

It is convenient for transparent calculations to use the notation defined by

$$(\tilde{T}^a)_{\alpha\alpha} \equiv \begin{cases} (T^{\tilde{a}})_{\alpha\alpha} & \text{for } a=\tilde{a} \quad (\text{not summed over } \alpha),\\ 0 & \text{for } a\neq\tilde{a}.\end{cases} \tag{4.9}$$

By taking account of the above discussions, the equation (4.1) becomes

$$\begin{aligned}&\mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak},\hat{A}_k^a,\hat{\Pi}_\psi,\hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\text{axial}}\\ &= \sum_{\alpha=1}^D\int \mathcal{D}A_\mu^a\mathcal{D}\bar{\psi}\mathcal{D}\psi\ \det(\partial_3)\prod_{x,a}\delta(A_3^a)\ \exp\left\{ig\int_0^\beta d\tau A_\tau^a(\tau,\mathbf{x}_0)(\tilde{T}^a)_{\alpha\alpha}\right\}\\ &\quad\times\exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{\frac{-1}{4}F_{\mu\nu}^aF_{\mu\nu}^a - \bar{\psi}i\gamma_\mu D_\mu\psi\Big\},\end{aligned} \tag{4.10}$$

where we have removed the dash $\prime$ from all the fields. Thus, our aim is now to find the operator $\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})$ which satisfies the above relation.

We propose the operator $\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})$ satisfying the Eq.(4.10) as

$$\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi}) = \sum_{\alpha=1}^{D} \exp\Big\{ g \int_0^\beta d\tau \int d^3\mathbf{x} (\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \Big((\partial_3)^{-2} \big((D_k \Pi_A^k)^a + J_F^{a0} \big) \Big) \Big\}, \tag{4.11}$$

where $(\tilde{T}^a)_{\alpha\alpha}$ is defined in Eq.(4.9). It should be emphasized that the above operator is described by only the physical operators but not A_τ. This is a main result of our paper. The above form of the operator might be guessed from Eqs.(2.5) and (2.9), but the proof that Eq.(4.11) leads to the relation (4.10) seems to be far from trivial, as we will see below.

For the operator (4.11), the path integral representation for the left-hand-side of Eq.(4.10) is

$$\begin{aligned}
&\mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})} \right)_{\text{axial}} = \sum_{\alpha=1}^{D} \int \mathcal{D}\Pi_A^{ak} \mathcal{D} A_k^a \mathcal{D}\Pi_\psi \mathcal{D}\psi \\
&\times \exp \int_0^\beta d\tau \int d^3\mathbf{x} \Big\{ g(\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \Big((\partial_3)^{-2} \big((D_k \Pi_A^k)^a + J_F^{a0} \big) \Big) \\
&+ i\Pi_A^{ak} \dot{A}_k^a + i\Pi_\psi \dot{\psi} - \mathcal{H}_{\text{axial}} \Big\},
\end{aligned} \tag{4.12}$$

where the Hamiltonian $\mathcal{H}_{\text{axial}}$ is given by Eq.(2.12). Since the exponent of the right-hand-side in Eq.(4.12) is quadratic with respect Π_A^{ak}, whose explicit calculations and forms are given in the Appendix A.3, we can perform the Gaussian integration $\mathcal{D}\Pi_A^{ak}$, and we obtain that

$$\begin{aligned}
&\mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})} \right)_{\text{axial}} \\
&= \sum_{\alpha=1}^{D} \int \mathcal{D} A_k^a \mathcal{D}\Pi_\psi \mathcal{D}\psi\ \det{}^{-\frac{1}{2}}(M^{-1}) \exp \int_0^\beta d\tau \int d^3\mathbf{x} \\
&\times \Big\{ -\frac{1}{2} \dot{A}_k^a \dot{A}_l^a - \frac{1}{2} (D_k \dot{A}_k)^a (\mathcal{D}^{-1} D_l \dot{A}_l)^a - i \dot{A}_k^a (D_k \mathcal{D}^{-1} J_F^0)^a + \frac{1}{2} J_F^{a0} (\mathcal{D}^{-1} J_F^0)^a \\
&- \frac{1}{4} F_{kl}^a F_{kl}^a - \frac{1}{2} \partial_3 A_k^a \partial_3 A_k^a + i\Pi_\psi \dot{\psi} + \bar{\psi} i\gamma^k D_k \psi + \bar{\psi} i\gamma^3 \partial_3 \psi \\
&+ g(\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) (\mathcal{D}^{-1} J_F^0)^a + i g(\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) (\mathcal{D}^{-1} D_k \dot{A}_k)^a \\
&- \frac{g^2}{2} \big((\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \big) \big((\partial_3)^{-2} (\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \big) \\
&+ \frac{g^2}{2} \big((\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \big) \big((\mathcal{D}^{-1})^{ab} (\tilde{T}^b)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \big) \Big\}.
\end{aligned} \tag{4.13}$$

We next insert the Gaussian integral,

$$
\begin{aligned}
1 = & \int \mathcal{D}A_\tau^a \, \det{}^{\frac{1}{2}}(\mathcal{D}) \, \exp \int_0^\beta d\tau \int d^3\mathbf{x} \\
& \Big\{ -\frac{1}{2} \Big(i A_\tau^a - (\mathcal{D}^{-1})^{ac} \big(i (D_k \dot{A}_k)^c + J_F^{c0} + g(\tilde{T}^c)_{\alpha\alpha} \delta^{(3)}(\mathbf{x} - \mathbf{x}_0) \big) \Big) \\
& \times (\mathcal{D})^{ad} \Big(i A_\tau^d - (\mathcal{D}^{-1})^{de} \big(i (D_l \dot{A}_l)^e + J_F^{e0} + g(\tilde{T}^e)_{\alpha\alpha} \delta^{(3)}(\mathbf{x} - \mathbf{x}_0) \big) \Big) \Big\}. \qquad (4.14)
\end{aligned}
$$

into Eq.(4.13) in order to recover the A_τ^a degree of freedom. Let us note that we add new terms of the point color charge densities $g(\tilde{T}^c)_{\alpha\alpha}\delta(\mathbf{x} - \mathbf{x}_0)$, compared with Eq.(3.8). If we expand the exponent of Eq.(4.14), which is given by Eq.(A.16) in the Appendix A, we see that some of the terms in the exponent of Eq.(4.13) are cancelled with those in Eq.(4.14). Then, we have

$$
\begin{aligned}
& \mathrm{Tr} \left(\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi}) \, \mathrm{e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})} \right)_{\text{axial}} \\
& = \sum_{\alpha=1}^{D} \int \mathcal{D}A_\tau^a \mathcal{D}A_k^a \mathcal{D}\Pi_\psi \mathcal{D}\psi \, \det{}^{-\frac{1}{2}}(M^{-1}) \det{}^{\frac{1}{2}}(\mathcal{D}) \, \exp \int_0^\beta d\tau \int d^3\mathbf{x} \\
& \Big\{ -\frac{1}{2} \dot{A}_k^a \dot{A}_k^a - \frac{1}{4} F_{kl}^a F_{kl}^a - \frac{1}{2} \partial_3 A_k^a \partial_3 A_k^a + i \Pi_\psi \dot{\psi} + \bar{\psi} i \gamma^k D_k \psi + \bar{\psi} i \gamma^3 \partial_3 \psi + i A_\tau^a J_F^{a0} \\
& - \frac{1}{2} (D_k A_\tau)^a (D_k A_\tau)^a - \frac{1}{2} \partial_3 A_\tau^a \partial_3 A_\tau^a + \dot{A}_k^a (D_k A_\tau)^a \\
& + i g A_\tau^a (\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x} - \mathbf{x}_0) \\
& - \frac{g^2}{2} \big((\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x} - \mathbf{x}_0) \big) \big((\partial_3)^{-2} (\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x} - \mathbf{x}_0) \big) \Big\}. \qquad (4.15)
\end{aligned}
$$

We note that A_τ^a is recovered as the auxiliary field through the Gaussian integral (4.14). The fourth line of Eq.(4.15) is just what we wanted and corresponds to the Polyakov loop in Eq.(4.10). Since the term comes from the cross terms in the Gaussian integral (4.14), it turns out to be difficult to get the Polyakov loop in the original path-ordered form of Eq.(4.1) from the above procedure. This is the reason why we rewrite the Polyakov loop of Eq.(4.1) into the abelian form given in Eq.(4.10).

We finally restore the A_3^a degree of freedom by Eq.(3.10) and the terms in the exponent (4.15) is summarized into the covariant form of the Lagrangian. The determinants are evaluated as before like Eq.(3.11). Then, we arrive at

$$
\begin{aligned}
& \mathrm{Tr} \left(\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi}) \, \mathrm{e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})} \right)_{\text{axial}} \\
= & \sum_{\alpha=1}^{D} \int \mathcal{D}A_\mu^a \mathcal{D}\bar{\psi} \mathcal{D}\psi \, \det(\partial_3) \prod_{x,a} \delta(A_3^a) \, \exp \Big\{ i g \int_0^\beta d\tau A_\tau^a(\tau, \mathbf{x}_0) (\tilde{T}^a)_{\alpha\alpha} \Big\} \\
& \times \exp \int_0^\beta d\tau \int d^3\mathbf{x} \Big\{ \frac{-1}{4} F_{\mu\nu}^a F_{\mu\nu}^a - \bar{\psi} i \gamma_\mu D_\mu \psi \Big\}. \qquad (4.16)
\end{aligned}
$$

Hence, we have proved Eq.(4.10) for the operator $\hat{\mathcal{O}}$ defined by Eq.(4.11). One can come back to the equation (4.1) by the inverse gauge transformations for Eqs.(4.2) $\sim$ (4.4). One can say that we prove the equation (3.3) in the axial gauge for the operator $\mathcal{O}$ defined by Eq.(4.11).

Before closing this section, let us discuss the last term in Eq.(4.15), which is the divergent self-energy of the point color charge densities due to introducing the Polyakov loop operator (4.11). If we define a new operator $\hat{\mathcal{O}}_{\rm new}$ in such a way that we subtract the self-energy part from the beginning by the counter term, $\delta^a_{\rm self}$,

$$\hat{\mathcal{O}}_{\rm new} \equiv \sum_{\alpha=1}^{D} \exp\left\{ g \int_0^\beta d\tau \int d^3\mathbf{x} (\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \Big((\partial_3)^{-2} \big((D_k \Pi^k_A)^a + J^{a0}_F + \delta^a_{\rm self} \big) \Big) \right\}, \quad (4.17)$$

where

$$\delta^a_{\rm self} \equiv \frac{g}{2} (\tilde{T}^a)_{\alpha\alpha} \delta^{(3)}(\mathbf{x}-\mathbf{x}_0), \quad (4.18)$$

then, the divergent term does not appear in the expression (4.15). We note that the term subtracted by the counter term, $(\partial_3)^{-2}\delta^a_{self}$, obviously does not depend on the field.

5 Hamiltonian in Coulomb Gauge

In this section with the following two subsections we reformulate the analyses done in the previous sections, 2, 3 and 4 for the case of the Coulomb gauge.

The Coulomb gauge is given by

$$\partial_i A^{ai} = 0 \qquad (i = 1, 2, 3). \quad (5.1)$$

One of the components, say, A^a_3 is eliminated by Eq.(5.1) like

$$A^a_3 = -(\partial_3)^{-1} \partial_k A^a_k. \quad (5.2)$$

We impose the Coulomb gauge on the $\nu = 0$ component of the equation of motion (2.3) and formally solve it for A^a_0. Then, we have

$$A^a_0 = (\mathcal{D}^{-1})^{ab} \Big((C_k \dot{A}_k)^b + J^{b0}_F \Big) \qquad (k = 1, 2). \quad (5.3)$$

The operator $(\mathcal{D}^{-1})^{ab}$ is defined as the inverse of $(\mathcal{D})^{ab} \equiv (D_i^2)^{ab}$. We find it useful to leave A^a_3 in the covariant derivative $(D_3)^{ab}$ as it is in order to perform calculations as clear as possible, keeping the equation (5.2) in mind. $(C_k)^{ab}$ in Eq.(5.3) is defined by

$$(C_k)^{ab} \equiv (D_k)^{ab} - (D_3)^{ab} (\partial_3)^{-1} \partial_k. \quad (5.4)$$

The partial integration with respect to $(C_k)^{ab}$ yields a new operator, accompanying a minus sign, given by

$$(\tilde{C}_k)^{ab} \equiv (D_k)^{ab} - \partial_k (\partial_3)^{-1} (D_3)^{ab} \quad (5.5)$$

and vice versa.

Likewise the case for the axial gauge, we eliminate A^a_0 by Eq. (5.3) and A^a_3 by Eq.(5.2) in the Lagrangian (2.1). Then, we obtain the Lagrangian $\mathcal{L}(A_k, \psi)_{\rm Coul}$ written in terms of only

physical degrees of freedom, whose explicit form is presented by Eq.(B.5) in the Appendix B. Then, the canonical momentum for A_k^a is given by

$$\Pi_A^{ak} \equiv \frac{\partial \mathcal{L}(A_k,\psi)_{\rm Coul}}{\partial \dot{A}_k^a} = (N)_{kl}^{ab}\dot{A}_l^b - (\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a, \tag{5.6}$$

where

$$(N)_{kl}^{ab} \equiv (\delta_{kl} + \partial_k(\partial_3)^{-2}\partial_l)\delta^{ab} - (\tilde{C}_k\mathcal{D}^{-1}C_l)^{ab}. \tag{5.7}$$

We obtain from Eq.(5.6) that

$$\dot{A}_k^a = (N^{-1})_{kl}^{ab}\Big(\Pi_A^{bl} + (\tilde{C}_l\mathcal{D}^{-1}J_F^0)^b\Big), \tag{5.8}$$

where $(N^{-1})_{kl}^{ab}$ is defined as the inverse of $(N)_{kl}^{ab}$ and its explicit form is

$$(N^{-1})_{kl}^{ab} = (\delta_{kl} - \partial_k\Delta^{-1}\partial_l)\delta^{ab} + \big(\tilde{C}'_k\tilde{\Delta}^{-1}C'_l\big)^{ab}, \tag{5.9}$$

where

$$\begin{aligned}
(\tilde{C}'_k)^{ab} &= (D_k)^{ab} - \partial_k\Delta^{-1}\partial\cdot(D)^{ab} && (5.10)\\
(C'_k)^{ab} &= (D_k)^{ab} - (D)^{ab}\cdot\partial\Delta^{-1}\partial_k, && (5.11)\\
\tilde{\Delta}^{-1} &\equiv \big(D\cdot\partial\Delta^{-1}\partial\cdot D\big)^{-1}, && (5.12)\\
\Delta^{-1} &\equiv (\partial_i^2)^{-1}. && (5.13)
\end{aligned}$$

We have also used the notation, $\partial\cdot D \equiv \partial_i D_i$. Let us note that $\partial\cdot D = D\cdot\partial$ in the Coulomb gauge. We present the relations among the operators, (5.4), (5.5), (5.10), (5.11) and the proof of Eq.(5.9) in the Appendix B. As for the fermion, we take the right derivative with respect to the Grassmann variable, so that we have

$$\Pi_\psi \equiv \frac{\partial \mathcal{L}(A_k,\psi)_{\rm Coul}}{\partial\dot{\psi}} = \bar{\psi}i\gamma^0. \tag{5.14}$$

Then, the Hamiltonian is obtained as

$$\begin{aligned}
\mathcal{H}_{\rm Coul} &= \Pi_A^{ak}\dot{A}_k^a + \Pi_\psi\dot{\psi} - \mathcal{L}(A_k,\psi)_{\rm Coul}\\
&= \frac{1}{2}\Pi_A^{ak}(N_{kl}^{-1}\Pi_A^l)^a + \Pi_A^{ak}(N_{kl}^{-1}\tilde{C}_l\mathcal{D}^{-1}J_F^0)^a\\
&\quad + \frac{1}{2}(\tilde{C}_k\mathcal{D}^{-1}J_F^0)(N_{kl}^{-1}\tilde{C}_l\mathcal{D}^{-1}J_F^0)^a - \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a\\
&\quad + \frac{1}{4}F_{kl}^aF_{kl}^a + \frac{1}{2}F_{3k}^aF_{3k}^a - \bar{\psi}i\gamma^kD_k\psi - \bar{\psi}i\gamma^3D_3\psi. && (5.15)
\end{aligned}$$

This Hamiltonian is written in terms of only physical degrees of freedom as it should. If we compare the Hamiltonian (5.15) with that in the axial gauge (2.12), we observe the correspondences among the operators in the Hamiltonians such as $M^{-1}\leftrightarrow N^{-1}, D_k\leftrightarrow\tilde{C}_k$ even though their explicit forms are quite different. The similarity between them may be the consequence of the fact that in the both cases the physical degrees of freedom are A_k, Π_A^{ak}, ψ and Π_ψ alone. The Hamiltonian (5.15) can be recast into another form, which is presented by Eq.(B.17) in the Appendix B, and the self-energy of the point color charge densities of fermion in the Coulomb gauge becomes manifest. We shall use the Hamiltonian (5.15) in the discussions below.

5.1 Trace Formula in Coulomb Gauge at Finite Temperature

In this section we would like to repeat the same analyses as the case of the axial gauge done in the section 3. We first show that the Hamiltonian (5.15) can reproduce the well-known path integral representation for the partition function in the Coulomb gauge at finite temperature,

$$\begin{aligned}\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} &= \int \mathcal{D}A_\mu^a \mathcal{D}\bar{\psi}\mathcal{D}\psi\ \det(\partial_i D_i)\prod_{x,a}\delta(\partial_i A_i^a) \\ &\quad\times \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{\frac{-1}{4}F_{\mu\nu}^a F_{\mu\nu}^a - \bar{\psi}i\gamma_\mu D_\mu\psi\Big\}. \qquad (5.16)\end{aligned}$$

The discussions here may be too pedagogical again, but is helpful for the discussions in the next subsection. The path integral representation for the left-hand-side of Eq.(5.16) is

$$\begin{aligned}&\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} \\ &= \int \mathcal{D}\Pi_A^{ak}\mathcal{D}A_k^a\mathcal{D}\Pi_\psi\mathcal{D}\psi\ \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{i\Pi_A^{ak}\dot{A}_k^a + i\Pi_\psi\dot{\psi} - \mathcal{H}_{\mathrm{Coul}}\Big\}. \qquad (5.17)\end{aligned}$$

Since the exponent in Eq.(5.17) is the quadratic with respect to Π_A^{ak}, one can perform the Gaussian integration. The detailed expression for the quadratic form is given by Eq.(B.18) in the Appendix B. Then, we have

$$\begin{aligned}&\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} = \int \mathcal{D}A_k^a\mathcal{D}\Pi_\psi\mathcal{D}\psi\ \det{}^{-\frac{1}{2}}(N^{-1})\ \exp\int_0^\beta d\tau\int d^3\mathbf{x} \\ &\times\Big\{-\frac{1}{2}\dot{A}_k^a\dot{A}_k^a - \frac{1}{2}\big((\partial_3)^{-1}\partial_k\dot{A}_k^a\big)\big((\partial_3)^{-1}\partial_l\dot{A}_l^a\big) + \frac{1}{2}\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}C_l\dot{A}_l)^a \\ &-i\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a + \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a \\ &-\frac{1}{4}F_{kl}^aF_{kl}^a - \frac{1}{2}F_{3k}^aF_{3k}^a + i\Pi_\psi\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3D_3\psi\Big\}. \qquad (5.18)\end{aligned}$$

We next consider the Gaussian integral given by

$$\begin{aligned}1 = \int \mathcal{D}A_\tau^a\ \det{}^{\frac{1}{2}}(\mathcal{D})\ \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{-\frac{1}{2}\Big(iA_\tau^a - (\mathcal{D}^{-1})^{ac}\big(i(C_k\dot{A}_k)^c + J_F^{c0}\big)\Big) \\ \times\ (\mathcal{D})^{ad}\Big(iA_\tau^d - (\mathcal{D}^{-1})^{de}\big(i(C_l\dot{A}_l)^e + J_F^{e0}\big)\Big)\Big\} \qquad (5.19)\end{aligned}$$

in order to restore the A_τ degree of freedom. Inserting the Gaussian integration (5.19) into Eq.(5.18), some of the terms in the exponents of Eqs.(5.18) and (5.19) are cancelled each other. We present the expansion of the exponent of Eq.(5.19) in Eq.(B.20) in the Appendix

B. Then, we have

$$\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}^k_A,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} = \int \mathcal{D}A^a_\tau \mathcal{D}A^a_k \mathcal{D}\Pi_\psi \mathcal{D}\psi \ \det{}^{-\frac{1}{2}}(N^{-1}) \ \det{}^{\frac{1}{2}}(\mathcal{D})$$
$$\times \exp\int_0^\beta d\tau \int d^3\mathbf{x}\Big\{-\frac{1}{2}\dot{A}^a_k\dot{A}^a_k - \frac{1}{2}\big((\partial_3)^{-1}\partial_k\dot{A}^a_k\big)\big((\partial_3)^{-1}\partial_l\dot{A}^a_l\big) - \frac{1}{4}F^a_{kl}F^a_{kl} - \frac{1}{2}F^a_{3k}F^a_{3k}$$
$$-\frac{1}{2}(D_iA_\tau)^a(D_iA_\tau)^a + \dot{A}^a_k(D_kA_\tau)^a - (D_3A_\tau)^a((\partial_3)^{-1}\partial_k\dot{A}^a_k)$$
$$+i\Pi_\psi\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3D_3\psi + iA^a_\tau J^{a0}_F\Big\}. \tag{5.20}$$

As in the case of the axial gauge, A_τ is recovered as the auxiliary field through the Gaussian integral (5.19). In order to restore the A^a_3 degree of freedom, let us consider the identity,

$$1 = \int \mathcal{D}A^a_3 \prod_{x,a}\delta(A^a_3 + (\partial_3)^{-1}\partial_kA^a_k) \ \Big(= \int \mathcal{D}A^a_3 \ \det(\partial_3)\prod_{x,a}\delta(\partial_iA^a_i)\Big). \tag{5.21}$$

Then, the exponent (5.20) is summarized into the covariant form of the Lagrangian thanks to the delta function in Eq.(5.21). And as shown in the Appendix C.2, the determinants are evaluated as

$$\det{}^{-\frac{1}{2}}(N^{-1})\det{}^{\frac{1}{2}}(\mathcal{D})\det(\partial_3) = \det(\partial_iD_i). \tag{5.22}$$

Hence, we obtain that

$$\mathrm{Tr}\left(\mathrm{e}^{-\beta H(\hat{\Pi}^k_A,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} = \int \mathcal{D}A^a_\mu \mathcal{D}\bar{\psi}\mathcal{D}\psi \ \det(\partial_iD_i)\prod_{x,a}\delta(\partial_iA^a_i)$$
$$\times \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{\frac{-1}{4}F^a_{\mu\nu}F^a_{\mu\nu} - \bar{\psi}i\gamma_\mu D_\mu\psi\Big\}, \tag{5.23}$$

where we have used $\mathcal{D}\Pi_\psi = \mathcal{D}\bar{\psi}$ and defined $\gamma_\tau \equiv -i\gamma^0$. We have finished to prove the equation (5.16).

5.2 Polyakov Loop in Coulomb Gauge

Let us proceed to discuss the Polyakov loop in the Coulomb gauge [7]. We no longer have the A_τ degree of freedom, from which the Polyakov loop is usually defined, in the operator formalism of the Coulomb gauge. Nevertheless, it is natural to expect that the operator written in terms of only physical degrees of freedom corresponding to the Polyakov loop should exist and to be defined.

[7] As a different approach from ours, a Polyakov loop in the temporal gauge has been discussed in Ref.[14] and treated as a Wilson line of the Hosotani mechanism[8] by exchanging the role of a spacial coordinate and the Euclidean time.

Our aim is to present the operator $\hat{\mathcal{O}}$ satisfying the equation (3.3) for the case of the Coulomb gauge, that is, it satisfies

$$\begin{aligned}
&\mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak},\hat{A}_k^a,\hat{\Pi}_\psi,\hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} \\
&= \sum_{\alpha=1}^{D}\int \mathcal{D}A_\mu^a\mathcal{D}\bar{\psi}\mathcal{D}\psi\ \det(\partial_i D_i)\prod_{x,a}\delta(\partial_i A_i^a)\ \exp\left\{ig\int_0^\beta d\tau A_\tau^a(\tau,\mathbf{x}_0)(\tilde{T}^a)_{\alpha\alpha}\right\} \\
&\times \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{\frac{-1}{4}F_{\mu\nu}^aF_{\mu\nu}^a-\bar{\psi}i\gamma_\mu D_\mu\psi\Big\},
\end{aligned}\tag{5.24}$$

where as in the case of the axial gauge the Polyakov loop is reduced to the $U(1)$ gauge sector of the $SU(N)$ gauge group by the gauge transformations (4.2) $\sim$ (4.4). Let us note again that $\hat{\mathcal{O}}$ should be written in terms of only physical degrees of freedom.

We shall show that

$$\hat{\mathcal{O}}\equiv\sum_{\alpha=1}^{D}\exp\left\{g\int_0^\beta d\tau\int d^3\mathbf{x}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\Big((\tilde{\Delta}^{-1})^{ab}\big((C'_k\Pi_A^k)^b+J_F^{b0}+\delta_{\mathrm{self}}^b\big)\Big)\right\}\tag{5.25}$$

satisfies the equation (5.24). Here we have inserted the counter term defined by

$$\delta_{\mathrm{self}}^b\equiv\frac{g}{2}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\tag{5.26}$$

in the definition (5.25). Note that the counter term (5.26) is the same form as the one for the axial gauge (4.18) and it does not depend on fields. δ_{self}^b cancels the divergent self-energy of the point color charge densities arising from introducing the operator (5.25), that is, the Polyakov loop operator as we will see below. It may be worth while pointing out that the term subtracted by the counter term is $(\tilde{\Delta}^{-1})^{ab}\delta_{self}^b$, so that, unlike the case for the axial gauge, it depends on the gauge field through the operator $\tilde{\Delta}^{-1}$. The above form of the operator (5.25) might be guessed from Eqs.(5.3) and (5.8), but it seems to be far from trivial that the operator actually satisfies the equation (5.24), as we will see below.

The path integral representation for the left-hand-side of Eq.(5.24) is

$$\begin{aligned}
&\mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak},\hat{A}_k^a,\hat{\Pi}_\psi,\hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} \\
&= \sum_{\alpha=1}^{D}\int \mathcal{D}\Pi_A^{ak}\mathcal{D}A_k^a\mathcal{D}\Pi_\psi\mathcal{D}\psi \\
&\times \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\Big((\tilde{\Delta}^{-1})^{ab}\big((C'_k\Pi_A^k)^b+J_F^{b0}+\delta_{\mathrm{self}}^b\big)\Big) \\
&+i\Pi_A^{ak}\dot{A}_k^a+i\Pi_\psi\dot{\psi}-\mathcal{H}_{\mathrm{Coul}}\Big\},
\end{aligned}\tag{5.27}$$

where the Hamiltonian $\mathcal{H}_{\mathrm{Coul}}$ is given by Eq.(5.15). The exponent in Eq.(5.27) is quadratic with respect to Π_A^{ak}, so that we can perform the integration $\mathcal{D}\Pi_A^{ak}$. The detailed calculations

and the explicit quadratic form for the exponent in Eq.(5.27) are given in the Appendix B.3. Then, we have

$$
\begin{aligned}
&\mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak},\hat{A}_k^a,\hat{\Pi}_\psi,\hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} = \sum_{\alpha=1}^{D}\int \mathcal{D}A_k^a\mathcal{D}\Pi_\psi\mathcal{D}\psi\ \det{}^{-\frac{1}{2}}(N^{-1}) \\
&\times \exp\int_0^\beta d\tau\int d^3\mathbf{x}\Big\{-\frac{1}{2}\dot{A}_k^a\dot{A}_k^a-\frac{1}{2}\big((\partial_3)^{-1}\partial_k\dot{A}_k^a\big)\big((\partial_3)^{-1}\partial_l\dot{A}_l^a\big)+\frac{1}{2}\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}C_l\dot{A}_l)^a \\
&-i\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a+\frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a-\frac{1}{4}F_{kl}^aF_{kl}^a-\frac{1}{2}F_{3k}^aF_{3k}^a \\
&+i\Pi_\psi\dot{\psi}+\bar{\psi}i\gamma^kD_k\psi+\bar{\psi}i\gamma^3D_3\psi \\
&+g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\tilde{\Delta}^{-1})^{ab}\delta^b_{\mathrm{self}} \\
&+g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}J_F^0)^a+ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}C_k\dot{A}_k)^a \\
&-\frac{g^2}{2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\tilde{\Delta}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) \\
&+\frac{g^2}{2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\mathcal{D}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big\}. \qquad (5.28)
\end{aligned}
$$

In order to recover the A_τ^a degree of freedom, we insert the Gaussian integral defined by

$$
\begin{aligned}
1 &= \int \mathcal{D}A_\tau^a\ \det{}^{\frac{1}{2}}(\mathcal{D})\ \exp\int_0^\beta d\tau\int d^3\mathbf{x} \\
&\times\Big\{-\frac{1}{2}\Big(iA_\tau^a-(\mathcal{D}^{-1})^{ac}\big(i(C_k\dot{A}_k)^c+J_F^{c0}+g(\tilde{T}^c)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big) \\
&\quad\times(\mathcal{D})^{ad}\Big(iA_\tau^d-(\mathcal{D}^{-1})^{de}\big(i(C_l\dot{A}_l)^e+J_F^{e0}+g(\tilde{T}^e)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big)\Big\}. \qquad (5.29)
\end{aligned}
$$

into Eq.(5.28). We have added new terms of the point color charge densities $g(\tilde{T}^c)_{\alpha\alpha}\delta(\mathbf{x}-\mathbf{x}_0)$, compared with Eq.(5.19). In Eq.(B.28) of the Appendix B, we explicitly present the expansion of the exponent of Eq.(5.29) and we find that some of the terms in the exponent of Eq.(5.29) are cancelled with those in Eq.(5.28). Then, we obtain that

$$
\begin{aligned}
&\mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak},\hat{A}_k^a,\hat{\Pi}_\psi,\hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k,\hat{A}_k,\hat{\Pi}_\psi,\hat{\psi})}\right)_{\mathrm{Coul}} \\
&=\sum_{\alpha=1}^{D}\int \mathcal{D}A_\tau^a\mathcal{D}A_k^a\mathcal{D}\Pi_\psi\mathcal{D}\psi\ \det{}^{-\frac{1}{2}}(N^{-1})\det{}^{\frac{1}{2}}(\mathcal{D})\ \exp\int_0^\beta d\tau\int d^3\mathbf{x} \\
&\times\Big\{-\frac{1}{2}\dot{A}_k^a\dot{A}_k^a-\frac{1}{2}\big((\partial_3)^{-1}\partial_k\dot{A}_k^a\big)\big((\partial_3)^{-1}\partial_l\dot{A}_l^a\big)-\frac{1}{4}F_{kl}^aF_{kl}^a-\frac{1}{2}F_{3k}^aF_{3k}^a \\
&-\frac{1}{2}(D_iA_\tau)^a(D_iA_\tau)^a+\dot{A}_k^a(D_kA_\tau)^a-(D_3A_\tau)^a((\partial_3)^{-1}\partial_k\dot{A}_k^a) \\
&+i\Pi_\psi\dot{\psi}+\bar{\psi}i\gamma^kD_k\psi+\bar{\psi}i\gamma^3D_3\psi+iA_\tau^aJ_F^{a0}+igA_\tau^a(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \\
&+g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\tilde{\Delta}^{-1})^{ab}\delta^b_{\mathrm{self}} \\
&-\frac{g^2}{2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\tilde{\Delta}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big\}. \qquad (5.30)
\end{aligned}
$$

The last two terms are found to be canceled with the choice of Eq.(5.26). The A_τ degree of freedom is recovered as the auxiliary field through the Gaussian integral (5.29). The last term in the fourth line of Eq.(5.30) just corresponds to the Polyakov loop in Eq.(5.24).

We finally restore the A_3^a degree of freedom by Eq.(5.21). As before, the exponent is summarized into the covariant form of the Lagrangian and the determinant is evaluated as Eq.(5.22). The result is

$$\mathrm{Tr}\left(\mathcal{O}(\hat{\Pi}_A^{ak}, \hat{A}_k^a, \hat{\Pi}_\psi, \hat{\psi})\ \mathrm{e}^{-\beta H(\hat{\Pi}_A^k, \hat{A}_k, \hat{\Pi}_\psi, \hat{\psi})}\right)_{\mathrm{Coul}} = \sum_{\alpha=1}^{D} \int \mathcal{D}A_\mu^a \mathcal{D}\bar{\psi}\mathcal{D}\psi\ \det(\partial_i D_i) \prod_{x,a} \delta(\partial_i A_i^a)$$
$$\times \exp\Big\{ig\int_0^\beta d\tau A_\tau^{\tilde{a}}(\tau, \mathbf{x}_0)(\tilde{T}^a)_{\alpha\alpha}\Big\}\exp\int_0^\beta d\tau \int d^3\mathbf{x}\Big\{\frac{-1}{4}F_{\mu\nu}^a F_{\mu\nu}^a - \bar{\psi}i\gamma_\mu D_\mu\psi\Big\}. \qquad (5.31)$$

Thus, we have proved the equation (5.24) and it implies that we have verified the equation (3.3) in the Coulomb gauge for the operator $\hat{\mathcal{O}}$ defined by Eq.(5.25).

It may be worth here mentioning the equivalence between the axial gauge and the Coulomb gauge. The Faddeev-Popov method guarantees the equivalence among different gauge choices in the path integral formalism even at finite temperature. Although the equivalence is believed to be true even in the operator formalism at finite temperature, the operator correspondence between operator formalisms with different gauge choices seems to be less obvious. Actually, our final results (4.16) and (5.31) show that the right-hand-sides of them are mutually transformed by the Faddeev-Popov manner. This fact implies that the results may prove indirectly the equivalence between the axial gauge and the Coulomb gauge in the operator formalism at finite temperature with the corresponding Polyakov loop operators.

6 Conclusions

In quantizing the non-abelian gauge theory with the non-covariant gauge fixing such as the axial and Coulomb gauge, any physical operator such as the Hamiltonian is written in terms of only physical degrees of freedom, by which the Hilbert space is spanned. A_0 is not dynamical variable, so that it disappears in the Hilbert space. Then, one may wonder how one should define the operator, written in terms of only physical degrees of freedom, corresponding to the Polyakov loop operator when one considers the theory in these gauge fixings.

In order to answer the question, we have first studied the canonical formulation of the non-abelian gauge theory in the axial and Coulomb gauge, in which the physical degrees of freedom are clarified. Namely, we emphasis on starting with the Lagrangian obtained by imposing the gauge condition and by eliminating A_0^a through the equation of motion, from which the canonical momentum for the gauge field is defined. The Hamiltonian obtained in this way is written in terms of only physical degrees of freedom as it should though it has the complicated form including the non-local operator.

We have constructed the operator corresponding to the Polyakov loop operator in terms of only physical degrees of freedom in the axial and Coulomb gauge. In order to confirm that the

defined operator is actually the standard Polyakov loop operator, we have evaluated the trace formula for the operator at finite temperature and showed that it can be rewritten into the covariant path integral form for the usual Polyakov loop operator with all the gauge degrees of freedom under the chosen gauge fixing.

We have encountered the divergent quantity in the process to obtain the covariant path integral form. It is the self-energy of the point color charge densities owing to introducing the operator $\hat{\mathcal{O}}$, which corresponds to the Polyakov loop. Introducing the Polyakov loop is essentially the same with considering the point color charge densities of fermion. This manifestly appears in the Hamiltonians, (A.6) and (B.17) in the Appendix. In the case of the axial gauge, the divergent self-energy does not depend on the field, while in the case of the Coulomb gauge it does. One can subtract the self-energy by introducing the counter term, $\delta^a_{\rm self}$ in the definition of the operator $\hat{\mathcal{O}}$, and the self-energy does not appear in the final expression of the covariant path integral form.

In proposing the operators (4.17) and (5.25), we have taken into account of the fact that the Polyakov loop operator has the same physical effect of introducing a color charge density. In fact, we have guessed the forms of the operators (4.17) and (5.25) from the second and third terms in the Hamiltonians (A.6) and (B.17). It is also very important and interesting to investigate the theoretical aspects of the non-abelian gauge theory at finite temperature based on the present work in addition to give the complete physical interpretation of the operator.

Acknowledgement

The authors thank Norisuke Sakai for raising the issue discussed in this paper. This work is supported in part by Grants-in-Aid for Scientific Research [No. 15K05055 and No. 25400260 (M.S.)] from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) in Japan.

A Axial gauge

In this Appendix we present some details of calculations and necessary formulae in order to derive the results in the text.

A.1 Hamiltonian in axial gauge

The Lagrangian $\mathcal{L}(A^a_k, \psi)_{\rm axial}$ is obtained by imposing the axial gauge $A^a_3 = 0$ and eliminating A^a_0 by using the constraint (2.5) in the Lagrangian (2.1). Straightforward calculations yield

$$\begin{aligned}\mathcal{L}(A^a_k, \psi)_{\rm axial} &= \frac{1}{2}\dot{A}^a_k (M_{kl}\dot{A}_l)^a - \dot{A}^a_k (D_k \mathcal{D}^{-1} J^0_F)^a + \frac{1}{2} J^{a0}_F (\mathcal{D}^{-1} J^0_F)^a \\ &\quad -\frac{1}{4}F^a_{kl}F^a_{kl} - \frac{1}{2}\partial_3 A^a_k \partial_3 A^a_k + \bar{\psi} i\gamma^0 \dot{\psi} + \bar{\psi} i\gamma^k D_k \psi + \bar{\psi} i\gamma^3 \partial_3 \psi, \end{aligned} \tag{A.1}$$

where we have used the definition for $(M)^{ab}_{kl}$, (2.8). In deriving (A.1), we have formally performed the partial integration with respect to D_k and $\mathcal{D}$. The partial integration with respect to D_k accompanies a minus sign, while that with respect to $\mathcal{D}$ does not. And we also note that the partial integration with respect to $(M)^{ab}_{kl}, (M^{-1})^{ab}_{kl}$ does not accompany a minus sign.

As mentioned in the section 2, the Hamiltonian (2.12),

$$\begin{aligned}
\mathcal{H}_{\rm axial} &= \Pi^{ak}_A \dot{A}^a_k + \Pi_\psi \dot{\psi} - \mathcal{L}(A_k, \psi)_{\rm axial} \\
&= \frac{1}{2}\Pi^{ak}_A (M^{-1}_{kl}\Pi^l_A)^a + \Pi^{ak}_A (M^{-1}_{kl} D_l \mathcal{D}^{-1} J^0_F)^a \\
&\quad + \frac{1}{2}(D_k \mathcal{D}^{-1} J^0_F)^a (M^{-1}_{kl} D_l \mathcal{D}^{-1} J^0_F)^a - \frac{1}{2} J^{a0}_F (\mathcal{D}^{-1} J^0_F)^a \\
&\quad + \frac{1}{4} F^a_{kl} F^a_{kl} + \frac{1}{2}\partial_3 A^a_k \partial_3 A^a_k - \bar{\psi} i\gamma^k D_k \psi - \bar{\psi} i\gamma^3 \partial_3 \psi
\end{aligned} \tag{A.2}$$

can be rewritten into another form, which is obtained by using the explicit form of $(M^{-1})^{ab}_{kl}$ for the second and the third term in Eq.(A.2) after the partial integration with respect to M^{-1}_{kl}, D_k and $\mathcal{D}^{-1}$. These two terms are written as

$$\begin{aligned}
&-(\mathcal{D}^{-1} D_l M^{-1}_{lk} \Pi^k_A)^a J^{a0}_F - \frac{1}{2} J^{a0}_F (\mathcal{D}^{-1} D_k M^{-1}_{kl} D_l \mathcal{D}^{-1} J^0_F)^a \\
&= \Pi^{ak}_A \Big(D_k (\partial_3)^{-2} J^0_F \Big)^a - \frac{1}{2} J^{a0}_F \Big(\big((\partial_3)^{-2} \delta^{ab} - (\mathcal{D}^{-1})^{ab} \big) J^{b0}_F \Big),
\end{aligned} \tag{A.3}$$

where we have used the relations,

$$(\mathcal{D}^{-1} D_l M^{-1}_{lk})^{ab} = (\partial_3)^{-2} (D_k)^{ab} \tag{A.4}$$

and

$$\Big(\mathcal{D}^{-1} D_k M^{-1}_{kl} D_l \mathcal{D}^{-1} \Big)^{ab} = (\partial_3)^{-2} \delta^{ab} - (\mathcal{D}^{-1})^{ab}, \tag{A.5}$$

which can be easily shown by using the explicit form of $(M^{-1})^{ab}_{kl}$ given by Eq.(2.10). We have finally performed the partial integration with respect to D_k and $(\partial_3)^{-2}$ in order to have the result (A.3). Then, we obtain that

$$\begin{aligned}
\mathcal{H}_{\rm axial} &= \frac{1}{2}\Pi^{ak}_A (M^{-1}_{kl}\Pi^l_A)^a + \Pi^{ak}_A (D_k (\partial_3)^{-2} J^0_F)^a - \frac{1}{2} J^{a0}_F (\partial_3)^{-2} J^{a0}_F \\
&\quad + \frac{1}{4} F^a_{kl} F^a_{kl} + \frac{1}{2}\partial_3 A^a_k \partial_3 A^a_k - \bar{\psi} i\gamma^k D_k \psi - \bar{\psi} i\gamma^3 \partial_3 \psi.
\end{aligned} \tag{A.6}$$

Let us note that the third term in the first line in Eq.(A.6) is well-known to be the self-energy of the point color charge densities of fermion in the axial gauge. The self-energy manifestly appears in this form of the Hamiltonian.

A.2 Trace formula in axial gauge

In order to perform the integration $\mathcal{D}\Pi_A^{ak}$, we complete the square with respect to Π_A^{ak} in the exponent of Eq. (3.5),

$$\begin{aligned}
& i\Pi_A^{ak}\dot{A}_k^a + i\Pi_\psi\dot{\psi} - \mathcal{H}_{\rm axial} \\
= & -\frac{1}{2}\Big(\Pi_A^{ak} - i(M_{kl}\dot{A}_l)^a + (D_k\mathcal{D}^{-1}J_F^0)^a\Big)(M^{-1})_{km}^{ab}\Big(\Pi_A^{bm} - i(M_{mn}\dot{A}_n)^b + (D_m\mathcal{D}^{-1}J_F^0)^b\Big) \\
& -\frac{1}{2}\dot{A}_k^a(M_{kl}\dot{A}_l)^a - i\dot{A}_k^a(D_k\mathcal{D}^{-1}J_F^0)^a + \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a - \frac{1}{4}F_{kl}^aF_{kl}^a - \frac{1}{2}\partial_3A_k^a\partial_3A_k^a \\
& +i\Pi_\psi\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3\partial_3\psi. \qquad \text{(A.7)}
\end{aligned}$$

The first term in the third line of Eq.(A.7) can be written as

$$-\frac{1}{2}\dot{A}_k^a(M_{kl}\dot{A}_l)^a = -\frac{1}{2}\dot{A}_k^a\dot{A}_k^a - \frac{1}{2}(D_k\dot{A}_k)^a(\mathcal{D}^{-1}D_l\dot{A}_l)^a \qquad \text{(A.8)}$$

by using the explicit form for $(M)_{kl}^{ab}$, (2.8) and by performing the partial integration for D_k. Then, we obtain the equation (3.7) in the text. If we expand the exponent of the Gaussian integral (3.8), we have, after performing the partial integration with respect to $\mathcal{D}$,

$$\begin{aligned}
& -\frac{1}{2}\Big(iA_\tau^a - (\mathcal{D}^{-1})^{ac}\big(i(D_k\dot{A}_k)^c + J_F^{c0}\big)\Big)(\mathcal{D})^{ad}\Big(iA_\tau^d - (\mathcal{D}^{-1})^{de}\big(i(D_l\dot{A}_l)^e + J_F^{e0}\big)\Big) \\
= & -\frac{1}{2}(D_kA_\tau)^a(D_kA_\tau)^a - \frac{1}{2}\partial_3A_\tau^a\partial_3A_\tau^a + iA_\tau^aJ_F^{a0} + \dot{A}_k^a(D_kA_\tau)^a \\
& +\frac{1}{2}(D_k\dot{A}_k)^a(\mathcal{D}^{-1}D_l\dot{A}_l)^a + i\dot{A}_k^a(D_k\mathcal{D}^{-1}J_F^0)^a - \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a. \qquad \text{(A.9)}
\end{aligned}$$

We observe that some of the terms in the second line of the right-hand-side of Eq. (A.7) and the right-hand-side of Eq.(A.9) are canceled each other to yield the result (3.9) in the text.

A.3 Polyakov loop in axial gauge

In order to perform the integration $\mathcal{D}\Pi_A^{ak}$, we need to complete the square with respect to Π_A^{ak} in the exponent of Eq.(4.12), which are given by

$$\begin{aligned}
& -\frac{1}{2}\Big(\Pi_A^{ak} - (M)_{kl}^{ab}\big(i\dot{A}_l^b - g(D_l)^{bd}(\partial_3)^{-2}(\tilde{T}^d)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) + (D_k\mathcal{D}^{-1}J_F^0)^b\Big)(M^{-1})_{km}^{ag} \\
& \times\Big(\Pi_A^{gm} - (M)_{mn}^{gc}\big(i\dot{A}_n^c - g(D_n)^{cf}(\partial_3)^{-2}(\tilde{T}^f)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) + (D_m\mathcal{D}^{-1}J_F^0)^g\Big). \qquad \text{(A.10)}
\end{aligned}$$

There are new terms containing the gauge coupling g due to introducing the operator (4.11), which do not appear in Eq.(A.7). According to the perfect square with respect to Π_A^{ak}, we should add the following terms,

$$\begin{aligned}
& -g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\partial_3)^{-2}(D_k\Pi_A^k)^a\big) + ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\partial_3)^{-2}D_kM_{kl}\dot{A}_l\big)^a \\
& +g(M)_{kl}^{ab}(D_l)^{bd}(\partial_3)^{-2}(\tilde{T}^d)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((M^{-1})_{km}^{ag}(D_m\mathcal{D}^{-1}J_F^0)^g\big) \\
& -\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\partial_3)^{-2}(D_kM_{kl}D_l)^{ab}(\partial_3)^{-2}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big), \qquad \text{(A.11)}
\end{aligned}$$

where we have performed the partial integration with respect to $(\partial_3)^{-2}, D_k$ and $(M)^{ab}_{kl}$. They can be rewritten into the simple forms by using the explicit form of M^{ab}_{kl} given by Eq.(2.8). It is easy to check that

$$\begin{aligned}(D_l M_{lk})^{ab} &= (D_l)^{ac}\big(\delta^{cb}\delta_{lk} - (D_l\mathcal{D}^{-1}D_k)^{cb}\big) = (D_k)^{ab} - (D_l^2\mathcal{D}^{-1}D_k)^{ab} \\ &= (D_k)^{ab} - \big((\mathcal{D}-(\partial_3)^2)\mathcal{D}^{-1}D_k\big)^{ab} = (\partial_3)^2(\mathcal{D}^{-1}D_k)^{ab} \end{aligned} \tag{A.12}$$

and

$$\begin{aligned}(D_k M_{kl} D_l)^{ab} &= (D_k M_{kl})^{ac}(D_l)^{cb} \overset{(A.12)}{=} (\partial_3)^2(\mathcal{D}^{-1}D_l)^{ac}(D_l)^{cb} \\ &= (\partial_3)^2\big(\mathcal{D}^{-1}(\mathcal{D}-(\partial_3)^2)\big)^{ab} = (D_k^2)^{ab} - (D_k^2\mathcal{D}^{-1}D_l^2)^{ab} \\ &= (\partial_3)^2\delta^{ab} - (\partial_3)^2(\mathcal{D}^{-1})^{ab}(\partial_3)^2. \end{aligned} \tag{A.13}$$

Then, the equation (A.11) becomes

$$\begin{aligned} & -g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\partial_3)^{-2}(D_k\Pi^k_A)^a\big) + ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\partial_3)^{-2}D_kM_{kl}\dot{A}_l\big)^a \\ & +g(M)^{ab}_{kl}(D_l)^{bd}(\partial_3)^{-2}(\tilde{T}^d)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((M^{-1})^{ag}_{km}(D_m\mathcal{D}^{-1}J^0_F)^g\big) \\ & -\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\partial_3)^{-2}(D_kM_{kl}D_l)^{ab}(\partial_3)^{-2}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) \\ = & -g\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big((\partial_3)^{-2}\big((D_k\Pi^k_A)^a + J^0_F\big)^a\Big) \\ & +g\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big(\mathcal{D}^{-1}J^0_F\big)^a + ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}D_k\dot{A}_k)^a \\ & -\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\partial_3)^{-2}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) \\ & +\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\mathcal{D}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big), \end{aligned} \tag{A.14}$$

where the first term, aside from the minus sign, in the right-hand-side is nothing but the exponent of the operator (4.11). Therefore, the exponent of the right-hand-side of Eq.(4.12) is written into the quadratic form with respect to Π^{ak}_A as

$$\begin{aligned} & g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\Big((\partial_3)^{-2}\big((D_k\Pi^k_A)^a + J^{a0}_F\big)\Big) + i\Pi^{ak}_A\dot{A}^a_k + i\Pi_\psi\dot{\psi} - \mathcal{H}_{\rm axial} \\ = & -\frac{1}{2}\Big(\Pi^{ak}_A - (M)^{ab}_{kl}\big(i\dot{A}^b_l - g(D_l)^{bd}(\partial_3)^{-2}(\tilde{T}^d)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) + (D_k\mathcal{D}^{-1}J^0_F)^b\Big)(M^{-1})^{ag}_{km} \\ & \times\Big(\Pi^{gm}_A - (M)^{gc}_{mn}\big(i\dot{A}^c_n - g(D_n)^{cf}(\partial_3)^{-2}(\tilde{T}^f)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) + (D_m\mathcal{D}^{-1}J^0_F)^g\Big) \\ & -\frac{1}{2}\dot{A}^a_k(M_{kl}\dot{A}_l)^a - i\dot{A}^a_k(D_k\mathcal{D}^{-1}J^0_F)^a + \frac{1}{2}J^{a0}_F(\mathcal{D}^{-1}J^0_F)^a - \frac{1}{4}F^a_{kl}F^a_{kl} - \frac{1}{2}\partial_3A^a_k\partial_3A^a_k \\ & +i\Pi_\psi\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3\partial_3\psi \\ & +g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}J^0_F)^a + ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}D_k\dot{A}_k)^a \\ & -\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\partial_3)^{-2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) \\ & +\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\mathcal{D}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big). \end{aligned} \tag{A.15}$$

We see that the equation (4.13) is obtained after the integration $\mathcal{D}\Pi_A^{ak}$. Let us note that the first term in the fourth line of Eq.(A.15) is rewritten by Eq.(A.8). If we expand the exponent of the Gaussian integral (4.14), we have

$$
\begin{aligned}
& -\frac{1}{2}\Big(iA_\tau^a - (\mathcal{D}^{-1})^{ac}\big(i(D_k\dot{A}_k)^c + J_F^{c0} + g(\tilde{T}^c)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big) \\
& \times(\mathcal{D})^{ad}\Big(iA_\tau^d - (\mathcal{D}^{-1})^{de}\big(i(D_l\dot{A}_l)^e + J_F^{e0} + g(\tilde{T}^e)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big) \\
= \; & -\frac{1}{2}(D_kA_\tau)^a(D_kA_\tau)^a - \frac{1}{2}\partial_3A_\tau^a\partial_3A_\tau^a + iA_\tau^aJ_F^{a0} + \dot{A}_k^a(D_kA_\tau)^a \\
& +\frac{1}{2}(D_k\dot{A}_k)^a(\mathcal{D}^{-1}D_l\dot{A}_l)^a + i\dot{A}_k^a(D_k\mathcal{D}^{-1}J_F^0)^a - \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a \\
& +igA_\tau^a(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \\
& -g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}J_F^0)^a - ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}D_k\dot{A}_k)^a \\
& -\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\mathcal{D}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big). \qquad \text{(A.16)}
\end{aligned}
$$

We find that some of the terms in Eqs.(A.15) and (A.16) are canceled each other, and we obtain the result (4.15) in the text. The fifth line in Eq.(A.16) corresponds to the Polyakov loop as it should. Let us note that the last term in Eq.(A.16) is canceled with the last term in Eq.(A.15). As for the term with the two delta functions, only the seventh line in Eq.(A.15) is left, which is the divergent self-energy of the point color charge densities owing to introducing the operator (4.11) as discussed in the text.

B Coulomb gauge

Let us present the formulae and relations in order to obtain the results of the section 5 in the text.

B.1 Hamiltonian in Coulomb gauge

Let us recall the operators defined in Eqs.(5.4) and (5.5), from which we obtain new operators given by Eqs.(5.10) and (5.11). These are proved by using $\partial_l^2 = \Delta - (\partial_3)^2$ like

$$
\begin{aligned}
(\tilde{C}_k')^{ab} &\equiv (\delta_{kl} - \partial_k\Delta^{-1}\partial_l)(\tilde{C}_l)^{ab} = (\delta_{kl} - \partial_k\Delta^{-1}\partial_l)\big((D_l)^{ab} - \partial_l(\partial_3)^{-1}(D_3)^{ab}\big) \\
&= (D_k)^{ab} - \partial_k\Delta^{-1}\partial\cdot(D)^{ab}, \qquad \text{(B.1)} \\
(C_k')^{ab} &\equiv (C_l)^{ab}(\delta_{lk} - \partial_l\Delta^{-1}\partial_k) = \big((D_l)^{ab} - (D_3)^{ab}(\partial_3)^{-1}\partial_l\big)(\delta_{lk} - \partial_l\Delta^{-1}\partial_k) \\
&= (D_k)^{ab} - (D)^{ab}\cdot\partial\Delta^{-1}\partial_k, \qquad \text{(B.2)}
\end{aligned}
$$

where we have used the notation $\partial_i(D_i)^{ab} = \partial\cdot(D)^{ab}$ which is also $(D)^{ab}\cdot\partial$ due to the Coulomb gauge, $\partial_iA^{ai} = 0$.

We also find the useful formulae given by

$$
\begin{aligned}
(C_k\tilde{C}'_k)^{ab} &= \big((D_k)^{ac} - (D_3)^{ac}(\partial_3)^{-1}\partial_k\big)\big((D_k)^{cb} - \partial_k\Delta^{-1}\partial\cdot(D)^{cb}\big) \\
&= (D_k^2)^{ab} + (D_3^2)^{ab} - (D_j\partial_j\Delta^{-1}\partial_i D_i)^{ab} \\
&= \mathcal{D}^{ab} - (\tilde{\Delta})^{ab}, \qquad \text{(B.3)} \\
(C'_k\tilde{C}_k)^{ab} &= \big((D_k)^{ac} - (D)^{ac}\cdot\partial\Delta^{-1}\partial_k\big)\big((D_k)^{cb} - \partial_l(\partial_3)^{-1}(D_3)^{ac}\big) \\
&= (D_k^2)^{ab} + (D_3^2)^{ab} - (D_i\partial_i\Delta^{-1}\partial_j D_j)^{ab} \\
&= \mathcal{D}^{ab} - (\tilde{\Delta})^{ab}, \qquad \text{(B.4)}
\end{aligned}
$$

where $\tilde{\Delta}$ is defined by $D_i\partial_i\Delta^{-1}\partial_j D_j$ whose inverse operator is given by Eq.(5.12).

The Lagrangian $\mathcal{L}(A_k^a,\psi)_{\rm Coul}$ is obtained by imposing the Coulomb gauge $\partial_i A^{ai}=0$ and eliminating A_0^a by using the constraint (5.3) in the Lagrangian (2.1), and after straightforward calculations we have

$$
\begin{aligned}
\mathcal{L}(A_k^a,\psi)_{\rm Coul} &= \frac{1}{2}\dot{A}_k^a(N_{kl}\dot{A}_l)^a - \dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a + \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a \\
&\quad -\frac{1}{4}F_{kl}^aF_{kl}^a - \frac{1}{2}F_{3k}^aF_{3k}^a + \bar{\psi}i\gamma^0\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3D_3\psi, \qquad \text{(B.5)}
\end{aligned}
$$

where we have used the definition (5.7). In deriving (B.5), we have formally performed the partial integration with respect to $\tilde{C}_k$ and $\mathcal{D}$. The partial integration with respect to $\tilde{C}_k$ gives a new operator C_k, accompanying a minus sign and vice versa. Incidentally, the partial integration with respect to $(N)_{kl}^{ab}, (N^{-1})_{kl}^{ab}$ does not accompany a minus sign.

Let us show that $(N^{-1})_{kl}^{ab}$ is given by Eq.(5.9). To this end, we first introduce a new operator defined by

$$
(N')_{km}^{ab} \equiv (\delta_{kl} - \partial_k\Delta^{-1}\partial_l)(N)_{lm}^{ab}. \qquad \text{(B.6)}
$$

$(N)_{km}^{ab}$ is defined by Eq.(5.7) in the text. Since it is easy to show that

$$
(\delta_{kl} - \partial_k\Delta^{-1}\partial_l)(\delta_{lm} + \partial_l(\partial_3)^{-2}\partial_m) = \delta_{km}, \qquad \text{(B.7)}
$$

the equation (B.6) becomes

$$
(N')_{km}^{ab} = \delta_{km}\delta^{ab} - (\tilde{C}'_k\mathcal{D}^{-1}C_m)^{ab}, \qquad \text{(B.8)}
$$

where we have used the definition (B.1). From Eq.(B.6), we obtain that

$$
(N^{-1})_{kl}^{ab} = (N^{-1\prime})_{km}^{ab}(\delta_{ml} - \partial_m\Delta^{-1}\partial_l). \qquad \text{(B.9)}
$$

The explicit form of $(N^{-1\prime})_{kl}^{ab}$ is easily found to be

$$
(N^{-1\prime})_{kl}^{ab} = \partial_{kl}\delta^{ab} + (\tilde{C}'_k\tilde{\Delta}^{-1}C_l)^{ab}. \qquad \text{(B.10)}
$$

This is, by using the relation (B.3), because

$$
\begin{aligned}
& (N')_{kl}^{ac}(N^{-1\prime})_{lm}^{cb} \\
&= \big(\delta_{kl}\delta^{ac} - (\tilde{C}'_k\mathcal{D}^{-1}C_l)^{ac}\big)\big(\partial_{lm}\delta^{cb} + (\tilde{C}'_l\tilde{\Delta}^{-1}C_m)^{cb}\big) \\
&= \delta_{km}\delta^{ab} + (\tilde{C}'_k\tilde{\Delta}^{-1}C_m)^{ab} - (\tilde{C}'_k\mathcal{D}^{-1}C_m)^{ab} - (\tilde{C}'_k\mathcal{D}^{-1}C_l\tilde{C}'_l\tilde{\Delta}^{-1}C_m)^{ab} \\
&= \delta_{km}\delta^{ab} + (\tilde{C}'_k\tilde{\Delta}^{-1}C_m)^{ab} - (\tilde{C}'_k\mathcal{D}^{-1}C_m)^{ab} - (\tilde{C}'_k\mathcal{D}^{-1}(\mathcal{D}-\tilde{\Delta})\tilde{\Delta}^{-1}C_m)^{ab} \\
&= \delta_{km}\delta^{ab}. \qquad \text{(B.11)}
\end{aligned}
$$

Then, the equation (B.9) is evaluated as

$$
\begin{aligned}
(N^{-1})^{ab}_{kl} &= \big(\delta_{km}\delta^{ab} + (\tilde{C}'_k\tilde{\Delta}^{-1}C_m)^{ab}\big)(\delta_{ml} - \partial_m\Delta^{-1}\partial_l) \\
&= (\delta_{kl} - \partial_k\Delta^{-1}\partial_l)\delta^{ab} + (\tilde{C}'_k\tilde{\Delta}^{-1}C'_l)^{ab},
\end{aligned} \tag{B.12}
$$

where we have used the definition (B.2) for C'_l. We have finished to prove the equation (5.9).

As in the case for the axial gauge, the Hamiltonian (5.15),

$$
\begin{aligned}
\mathcal{H}_{\rm Coul} &= \Pi^{ak}_A\dot{A}^a_k + \Pi_\psi\dot{\psi} - \mathcal{L}(A_k,\psi)_{\rm Coul} \\
&= \frac{1}{2}\Pi^{ak}_A(N^{-1}_{kl}\Pi^l_A)^a + \Pi^{ak}_A(N^{-1}_{kl}\tilde{C}_l\mathcal{D}^{-1}J^0_F)^a \\
&\quad + \frac{1}{2}(\tilde{C}_k\mathcal{D}^{-1}J^0_F)(N^{-1}_{kl}\tilde{C}_l\mathcal{D}^{-1}J^0_F)^a - \frac{1}{2}J^{a0}_F(\mathcal{D}^{-1}J^0_F)^a \\
&\quad + \frac{1}{4}F^a_{kl}F^a_{kl} + \frac{1}{2}F^a_{3k}F^a_{3k} - \bar{\psi}i\gamma^kD_k\psi - \bar{\psi}i\gamma^3D_3\psi
\end{aligned} \tag{B.13}
$$

can be rewritten into another form. The second and the third terms in Eq.(B.13) after the partial integration with respect to $N^{-1}_{kl}, \tilde{C}_k$ and $\mathcal{D}^{-1}$ are rewritten as

$$
\begin{aligned}
&-(\mathcal{D}^{-1}C_lN^{-1}_{lk}\Pi^k_A)^aJ^{a0}_F - \frac{1}{2}J^{0a}_F(\mathcal{D}^{-1}C_kN^{-1}_{kl}\tilde{C}_l\mathcal{D}^{-1}J^0_F)^a \\
&= \Pi^{ak}_A\big(\tilde{C}'_k\tilde{\Delta}^{-1}J^0_F\big)^a - \frac{1}{2}J^{a0}_F\Big(\big((\tilde{\Delta}^{-1})^{ab} - (\mathcal{D}^{-1})^{ab}\big)J^{b0}_F\Big),
\end{aligned} \tag{B.14}
$$

where we have used the relations,

$$
(\mathcal{D}^{-1}C_lN^{-1}_{lk})^{ab} = (\tilde{\Delta}^{-1}C'_k)^{ab} \tag{B.15}
$$

and

$$
\Big(\mathcal{D}^{-1}C_kN^{-1}_{kl}\tilde{C}_l\mathcal{D}^{-1}\Big)^{ab} = (\tilde{\Delta}^{-1})^{ab} - (\mathcal{D}^{-1})^{ab}. \tag{B.16}
$$

The above relations can be shown by using the equations (B.3), (B.4) and the explicit form of $(N^{-1})^{ab}_{kl}$ given by Eq.(B.12). We have finally performed the partial integration with respect to $\tilde{\Delta}^{-1}$ and C'_k in order to arrive at Eq.(B.14). Note that the partial integration with respect to C'_k yields a new operator $\tilde{C}'_k$, accompanying a minus sign and vice versa, while the partial integration of $\tilde{\Delta}^{-1}$ does not have a minus sign. Then, we obtain that

$$
\begin{aligned}
\mathcal{H}_{\rm Coul} &= \frac{1}{2}\Pi^{ak}_A(N^{-1}_{kl}\Pi^l_A)^a + \Pi^{ak}_A(\tilde{C}'_k\tilde{\Delta}^{-1}J^0_F)^a - \frac{1}{2}J^{a0}_F(\tilde{\Delta}^{-1}J^0_F)^a \\
&\quad + \frac{1}{4}F^a_{kl}F^a_{kl} + \frac{1}{2}F^a_{3k}F^a_{3k} - \bar{\psi}i\gamma^kD_k\psi - \bar{\psi}i\gamma^3D_3\psi.
\end{aligned} \tag{B.17}
$$

Let us note that the third term in the first line of Eq.(B.17) is well-known to be the self-energy of the point color charge densities of fermion in the Coulomb gauge. Unlike the case for the axial gauge, it depends on the gauge field through the operator $\tilde{\Delta}^{-1}$. We also point out the correspondence of the Hamiltonians (A.6) and (B.17) such as $M^{-1} \leftrightarrow N^{-1}, D_k \leftrightarrow \tilde{C}'_k$ and $(\partial_3)^{-2} \leftrightarrow \tilde{\Delta}^{-1}$, though the explicit forms for the operators are quite different.

B.2 Trace formula in Coulomb gauge

We complete the square with respect to Π_A^{ak} in the exponent of Eq.(5.17) in order to perform the integration $\mathcal{D}\Pi_A^{ak}$,

$$
\begin{aligned}
& i\Pi_A^{ak}\dot{A}_k^a + i\Pi_\psi\dot{\psi} - \mathcal{H}_{\rm Coul} \\
= \; & -\frac{1}{2}\Big(\Pi_A^{ak} - i(N_{kl}\dot{A}_l)^a + (\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a\Big)(N^{-1})_{km}^{ac}\Big(\Pi_A^{cm} - i(N_{mn}\dot{A}_n)^c + (\tilde{C}_m\mathcal{D}^{-1}J_F^0)^c\Big) \\
& -\frac{1}{2}\dot{A}_k^a(N_{kl}\dot{A}_l)^a - i\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a + \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a - \frac{1}{4}F_{kl}^aF_{kl}^a - \frac{1}{2}F_{3k}^aF_{3k}^a \\
& +i\Pi_\psi\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3D_3\psi.
\end{aligned}
\tag{B.18}
$$

The first term in the third line of Eq.(B.18) is rewritten as

$$
-\frac{1}{2}\dot{A}_k^a(N_{kl}\dot{A}_l)^a = -\frac{1}{2}\dot{A}_k^a\dot{A}_k^a - \frac{1}{2}\big((\partial_3)^{-1}\partial_k\dot{A}_k^a\big)\big((\partial_3)^{-1}\partial_l\dot{A}_l^a\big) + \frac{1}{2}\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}C_l\dot{A}_l)^a, \tag{B.19}
$$

where we have used the explicit form for $(N)_{kl}^{ab}$, Eq.(5.7) and performed the partial integration with respect to $(\partial_3)^{-1}$ and ∂_k.

The expansion of the exponent of the Gaussian integral (5.19) yields

$$
\begin{aligned}
& -\frac{1}{2}\Big(iA_\tau^a - (\mathcal{D}^{-1})^{ac}\big(i(C_k\dot{A}_k)^c + J_F^{c0}\big)\Big)(\mathcal{D})^{ad}\Big(iA_\tau^d - (\mathcal{D}^{-1})^{de}\big(i(C_l\dot{A}_l)^e + J_F^{e0}\big)\Big) \\
= \; & -\frac{1}{2}(D_iA_\tau)^a(D_iA_\tau)^a + iA_\tau^aJ_F^{a0} + \dot{A}_k^a(D_kA_\tau)^a - \frac{1}{2}\dot{A}_l^a(\tilde{C}_l\mathcal{D}^{-1}C_k\dot{A}_k)^a \\
& +i\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a - \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a
\end{aligned}
\tag{B.20}
$$

Let us note that the third and the fourth terms in Eq.(B.20) comes from one of the terms obtained by expanding the exponent of Eq.(B.20),

$$
-A_\tau^a\big(C_k\dot{A}_k\big)^a = \big(D_kA_\tau\big)^a\dot{A}_k^a - \big(D_3A_\tau\big)^a\big((\partial_3)^{-1}\partial_k\dot{A}_k^a\big), \tag{B.21}
$$

where we have used the equation (5.4) and the partial integration with respect to D_k has been done. We see that some of the terms in Eqs. (B.18) and (B.20) are canceled each other to yield the result (5.20) in the text.

B.3 Polyakov loop in Coulomb gauge

We need to complete the square with respect to Π_A^{ak} in the exponent of Eq.(5.27) in order to perform the integration $\mathcal{D}\Pi_A^{ak}$. It is given by

$$
\begin{aligned}
& -\frac{1}{2}\Big(\Pi_A^{ak} - (N)_{kl}^{ab}\big(i\dot{A}_l^b - g(\tilde{C}_l'\tilde{\Delta}^{-1})^{bd}(\tilde{T}^d)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) + (\tilde{C}_k\mathcal{D}^{-1}J_F^0)^b\Big)(N^{-1})_{km}^{ag} \\
& \times\Big(\Pi_A^{gm} - (N)_{mn}^{gc}\big(i\dot{A}_n^c - g(\tilde{C}_n'\tilde{\Delta}^{-1})^{cf}(\tilde{T}^f)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) + (\tilde{C}_m\mathcal{D}^{-1}J_F^0)^g\Big).
\end{aligned}
\tag{B.22}
$$

There appear new terms with the gauge coupling g because of introducing the operator (5.25), which we do not have in Eq.(B.18). According to the perfect square with respect to Π_A^{ak}, we should add the following terms,

$$
\begin{aligned}
&-g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big(\tilde{\Delta}^{-1}C'_k\Pi_A^k\big)^a + ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\tilde{\Delta}^{-1}C'_kN_{kl}\dot{A}_l)^a \\
&+g(N)^{ab}_{kl}(\tilde{C}'_l\tilde{\Delta}^{-1})^{bd}(\tilde{T}^d)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((N^{-1})^{ag}_{km}(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^g\big) \\
&-\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}(\partial_3)^{-2}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\tilde{\Delta}^{-1}C'_kN_{kl}\tilde{C}'_l\tilde{\Delta}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big), \qquad \text{(B.23)}
\end{aligned}
$$

where we have performed the partial integration with respect to $\tilde{\Delta}^{-1}, \tilde{C}'_k$ and $(N)^{ab}_{kl}$. They can be rewritten into simple forms by using the explicit form of N^{ab}_{kl}. With the help of Eqs.(B.2), (B.3), (B.4) and (B.7) we obtain that

$$
\begin{aligned}
(\tilde{\Delta}^{-1}C'_kN_{kl})^{ab} &= (\tilde{\Delta}^{-1}C'_k)^{ac}\big(\delta^{cb}(\delta_{kl}+\partial_k(\partial_3)^{-1}\partial_l) - (\tilde{C}_k\mathcal{D}^{-1}C_l)^{cb}\big) \\
&= (\tilde{\Delta}^{-1})^{ac}(C_l)^{cb} - (\tilde{\Delta}^{-1})^{ac}(C'_k\tilde{C}_k)^{cd}(\mathcal{D}^{-1}C_l)^{db} \\
&= (\tilde{\Delta}^{-1})^{ac}(C_l)^{cb} - (\tilde{\Delta}^{-1})^{ac}(\mathcal{D}-\tilde{\Delta})^{cd}(\mathcal{D}^{-1}C_l)^{db} \\
&= (\mathcal{D}^{-1}C_l)^{ab} \qquad \text{(B.24)}
\end{aligned}
$$

and

$$
\begin{aligned}
(\tilde{\Delta}^{-1}C'_kN_{kl}\tilde{C}'_l\tilde{\Delta}^{-1})^{ab} &= (\tilde{\Delta}^{-1}C'_kN_{kl})^{ac}(\tilde{C}'_l\tilde{\Delta}^{-1})^{cb} \overset{(B.24)}{=} (\mathcal{D}^{-1}C_l)^{ac}(\tilde{C}'_l\tilde{\Delta}^{-1})^{cb} \\
&= (\mathcal{D}^{-1})^{ac}(C_l\tilde{C}'_l)^{cd}(\tilde{\Delta}^{-1})^{db} = (\mathcal{D}^{-1})^{ac}(\mathcal{D}-\tilde{\Delta})^{cd}(\tilde{\Delta}^{-1})^{db} \\
&= (\tilde{\Delta}^{-1})^{ab} - (\mathcal{D}^{-1})^{ab}. \qquad \text{(B.25)}
\end{aligned}
$$

Then, we have

$$
\begin{aligned}
&-g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big(\tilde{\Delta}^{-1}C'_k\Pi_A^k\big)^a + ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\tilde{\Delta}^{-1}C'_kN_{kl}\dot{A}_l)^a \\
&+g(N)^{ab}_{kl}(\tilde{C}'_l\tilde{\Delta}^{-1})^{bd}(\tilde{T}^d)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((N^{-1})^{ag}_{km}(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^g\big)\big) \\
&-\frac{g^2}{2}\big((\tilde{T}^a)_{\alpha\alpha}(\partial_3)^{-2}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\big((\tilde{\Delta}^{-1}C'_kN_{kl}\tilde{C}'_l\tilde{\Delta}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) \\
= \;& -g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\Big((\tilde{\Delta}^{-1})^{ab}\big((C'_k\Pi_A^k)^a + J_F^{0b}\big)\Big) \\
&+g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}J_F^0)^a + ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}C_k\dot{A}_k^a) \\
&-\frac{g^2}{2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\tilde{\Delta}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) \\
&+\frac{g^2}{2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\mathcal{D}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big), \qquad \text{(B.26)}
\end{aligned}
$$

where the first term, aside from the minus sign, in the right-hand-side is the exponent of operator (5.25). Hence, the exponent of the right-hand-side of Eq.(5.27) is written into the

quadratic form with respect to Π_A^{ak} as

$$
\begin{aligned}
& g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\Big((\tilde{\Delta}^{-1})^{ab}\big((C'_k\Pi_A^k)^b + J_F^{b0} + \delta^b_{\rm self}\big)\Big) + i\Pi_A^{ak}\dot{A}_k^a + i\Pi_\psi\dot{\psi} - \mathcal{H}_{\rm Coul} \\
= & -\frac{1}{2}\Big(\Pi_A^{ak} - (N)^{ab}_{kl}\big(i\dot{A}_l^b - g(\tilde{C}'_l\tilde{\Delta}^{-1})^{bd}(\tilde{T}^d)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) + (\tilde{C}_k\mathcal{D}^{-1}J_F^0)^b\Big)(N^{-1})^{ag}_{km} \\
& \times\Big(\Pi_A^{gm} - (N)^{gc}_{mn}\big(i\dot{A}_n^c - g(\tilde{C}'_n\tilde{\Delta}^{-1})^{cf}(\tilde{T}^f)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) + (\tilde{C}_m\mathcal{D}^{-1}J_F^0)^g\Big) \\
& -\frac{1}{2}\dot{A}_k^a(N_{kl}\dot{A}_l)^a - i\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a + \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a - \frac{1}{4}F_{kl}^aF_{kl}^a - \frac{1}{2}F_{3k}^aF_{3k}^a \\
& +i\Pi_\psi\dot{\psi} + \bar{\psi}i\gamma^kD_k\psi + \bar{\psi}i\gamma^3D_3\psi \\
& +g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\tilde{\Delta}^{-1})^{ab}\delta^b_{\rm self} \\
& +g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}J_F^0)^a + ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}C_k\dot{A}_k)^a \\
& -\frac{g^2}{2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\tilde{\Delta}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big) \\
& +\frac{g^2}{2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\mathcal{D}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big). \qquad \text{(B.27)}
\end{aligned}
$$

We observe that the integration $\mathcal{D}\Pi_A^{ak}$ yields the equation (5.28). Let us note that the first term in the fourth line of Eq.(B.27) is rewritten by Eq.(B.19). As explained in the text, the counter term, the sixth line of Eq.(B.27), exactly cancels the divergent self-energy, the eighth line of Eq.Eq.(B.27).

If we expand the exponent of the Gaussian integral (5.29), we have

$$
\begin{aligned}
& -\frac{1}{2}\Big(iA_\tau^a - (\mathcal{D}^{-1})^{ac}\big(i(C_k\dot{A}_k)^c + J_F^{c0} + g(\tilde{T}^c)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big) \\
& \quad\times(\mathcal{D})^{ad}\Big(iA_\tau^d - (\mathcal{D}^{-1})^{de}\big(i(C_l\dot{A}_l)^e + J_F^{e0} + g(\tilde{T}^e)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big)\Big) \\
= & -\frac{1}{2}(D_iA_\tau)^a(D_iA_\tau)^a + iA_\tau^aJ_F^{a0} - A_\tau^a(C_k\dot{A}_k)^a \\
& -\frac{1}{2}\dot{A}_l^a(\tilde{C}_l\mathcal{D}^{-1}C_k\dot{A}_k)^a + i\dot{A}_k^a(\tilde{C}_k\mathcal{D}^{-1}J_F^0)^a - \frac{1}{2}J_F^{a0}(\mathcal{D}^{-1}J_F^0)^a \\
& +igA_\tau^a(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0) \\
& -g(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}J_F^0)^a - ig(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)(\mathcal{D}^{-1}C_k\dot{A}_k)^a \\
& -\frac{g^2}{2}(\tilde{T}^a)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big((\mathcal{D}^{-1})^{ab}(\tilde{T}^b)_{\alpha\alpha}\delta^{(3)}(\mathbf{x}-\mathbf{x}_0)\big). \qquad \text{(B.28)}
\end{aligned}
$$

The third term in the first line of the right-hand-side of Eq.(B.28) can be written by

$$
-A_\tau^a(C_k\dot{A}_k)^a = \dot{A}_k^a(D_kA_\tau)^a - (D_3A_\tau)^a((\partial_3)^{-1}\partial_k\dot{A}_k^a), \qquad \text{(B.29)}
$$

where we have performed the partial integration with respect to D_3. We observe that some of the terms in Eqs.(B.27) and (B.28) are canceled each other to result the equation (5.30) in the text.

C Evaluation of Determinants

C.1 Axial gauge

Let us prove the equation (3.11). We evaluate $\det(M)$ where M is defined by Eq.(2.8). To this end, we write M as

$$(M)^{ab}_{kl} = \begin{pmatrix} \delta^{ab} - (D_1\mathcal{D}^{-1}D_1)^{ab} & -(D_1\mathcal{D}^{-1}D_2)^{ab} \\ -(D_2\mathcal{D}^{-1}D_1)^{ab} & \delta^{ab} - (D_2\mathcal{D}^{-1}D_2)^{ab} \end{pmatrix} \qquad (a,b = 1 \sim N^2-1). \tag{C.1}$$

Introducing the relation given by

$$\det \begin{pmatrix} 0 & \mathbb{I}_{N^2-1} \\ \mathbb{I}_{N^2-1} & 0 \end{pmatrix} = (-1)^{N^2-1}, \tag{C.2}$$

where $\mathbb{I}_{N^2-1}$ is the $(N^2-1)\times(N^2-1)$ unit matrix, we have

$$\begin{aligned} (-1)^{N^2-1}\det(M) &= \det\left[\begin{pmatrix} 0 & \mathbb{I}_{N^2-1} \\ \mathbb{I}_{N^2-1} & 0 \end{pmatrix} M\right] \\ &= \det\begin{pmatrix} -D_2\mathcal{D}D_1 & \mathbb{I}_{N^2-1} - D_2\mathcal{D}^{-1}D_2 \\ \mathbb{I}_{N^2-1} - D_1\mathcal{D}^{-1}D_1 & -D_1\mathcal{D}^{-1}D_2 \end{pmatrix}. \end{aligned} \tag{C.3}$$

With help of the formula,

$$\det\begin{pmatrix} A & C \\ D & B \end{pmatrix} = \det\begin{pmatrix} A & 0 \\ D & \mathbb{I} \end{pmatrix}\begin{pmatrix} \mathbb{I} & A^{-1}C \\ 0 & B - DA^{-1}C \end{pmatrix} = \det(A)\times\det(B - DA^{-1}C), \tag{C.4}$$

The equation (C.3) becomes

$$\begin{aligned} & \det(-D_2\mathcal{D}D_1)\det\Big(-D_1\mathcal{D}^{-1}D_2 - (\mathbb{I}_{N^2-1} - D_1\mathcal{D}^{-1}D_1)(-D_2\mathcal{D}^{-1}D_1)^{-1}(\mathbb{I}_{N^2-1} - D_2\mathcal{D}^{-1}D_2)\Big) \\ = \;& \det\Big(D_2\mathcal{D}^{-1}(D_1)^2\mathcal{D}^{-1}D_2 - (D_2\mathcal{D}^{-1}D_1)(\mathbb{I}_{N^2-1} - D_1\mathcal{D}^{-1}D_1)(D_1^{-1}\mathcal{D}D_2^{-1})(\mathbb{I}_{N^2-1} - D_2\mathcal{D}^{-1}D_2)\Big) \\ = \;& \det\Big(D_2\mathcal{D}^{-1}(D_1)^2\mathcal{D}^{-1}D_2 - \mathbb{I}_{N^2-1} + D_2\mathcal{D}^{-1}D_2 + D_2\mathcal{D}^{-1}(D_1)^2D_2^{-1} - D_2\mathcal{D}^{-1}(D_1)^2\mathcal{D}^{-1}D_2\Big) \\ = \;& (-1)^{N^2-1}\det(D_2^{-1}D_2)\det\Big(\mathbb{I}_{N^2-1} - D_2\mathcal{D}^{-1}D_2 - D_2\mathcal{D}^{-1}(D_1)^2D_2^{-1}\Big) \\ = \;& (-1)^{N^2-1}\det\Big(D_2^{-1}\big(\mathbb{I}_{N^2-1} - D_2\mathcal{D}^{-1}D_2 - D_2\mathcal{D}^{-1}(D_1)^2D_2^{-1}\big)D_2\Big) \\ = \;& (-1)^{N^2-1}\det(\mathcal{D}^{-1}(\partial_3)^2), \end{aligned} \tag{C.5}$$

where we have used $\mathcal{D} = (D_1)^2 + (D_2)^2 + (\partial_3)^2$ in the last line in Eq.(C.5). Hence, we obtain that

$$\det(M) = \det\big(\mathcal{D}^{-1}(\partial_3)^2\big), \tag{C.6}$$

which is the equation (3.11) in the text.

C.2 Coulomb gauge

Instead of evaluating $\det(N)$ directly, let us consider the determinant defined by

$$\det\left(\begin{pmatrix} 0 & \mathbb{I}_{N^2-1} \\ \mathbb{I}_{N^2-1} & 0 \end{pmatrix} N'\right), \tag{C.7}$$

where N' is given by Eq.(B.8) and is written as

$$(N')^{ab}_{kl} = \begin{pmatrix} \mathbb{I}_{N^2-1} - \tilde{C}'_1\mathcal{D}^{-1}C_1 & -\tilde{C}'_1\mathcal{D}^{-1}C_2 \\ -\tilde{C}'_2\mathcal{D}^{-1}C_1 & \mathbb{I}_{N^2-1} - \tilde{C}'_2\mathcal{D}^{-1}C_2 \end{pmatrix}. \tag{C.8}$$

The equation (C.7) becomes, by using the formula (C.4),

$$\begin{aligned}
(C.7) &= \det\begin{pmatrix} -\tilde{C}'_2\mathcal{D}^{-1}C_1 & \mathbb{I}_{N^2-1} - \tilde{C}'_2\mathcal{D}^{-1}C_2 \\ \mathbb{I}_{N^2-1} - \tilde{C}'_1\mathcal{D}^{-1}C_1 & -\tilde{C}'_1\mathcal{D}^{-1}C_2 \end{pmatrix} \\
&= \det\Big(-\tilde{C}'_2\mathcal{D}^{-1}C_1\Big) \\
&\quad \times\det\Big(-\tilde{C}'_1\mathcal{D}^{-1}C_2 - (\mathbb{I}_{N^2-1} - \tilde{C}'_1\mathcal{D}^{-1}C_1)(-\tilde{C}'_2\mathcal{D}^{-1}C_1)^{-1}(\mathbb{I}_{N^2-1} - \tilde{C}'_2\mathcal{D}^{-1}C_2)\Big) \\
&= \det\Big(-\mathbb{I}_{N^2-1} + \tilde{C}'_2\mathcal{D}^{-1}C_2 + \tilde{C}'_2\mathcal{D}^{-1}C_1\tilde{C}'_1(\tilde{C}'_2)^{-1}\Big) \\
&= (-1)^{N^2-1}\,\det(\tilde{C}'^{-1}_2\tilde{C}'_2)\det\Big(\mathbb{I}_{N^2-1} - \tilde{C}'_2\mathcal{D}^{-1}C_2 - \tilde{C}'_2\mathcal{D}^{-1}C_1\tilde{C}'_1(\tilde{C}'_2)^{-1}\Big) \\
&= (-1)^{N^2-1}\,\det(\mathbb{I}_{N^2-1} - \mathcal{D}^{-1}C_2\tilde{C}'_2 - \mathcal{D}^{-1}C_1\tilde{C}'_1) \\
&= (-1)^{N^2-1}\,\det(\mathcal{D}^{-1})\det(\mathcal{D} - C_2\tilde{C}'_2 - C_1\tilde{C}'_1) \\
&= (-1)^{N^2-1}\,\det(\mathcal{D}^{-1})\det(\mathcal{D} - C_k\tilde{C}'_k) \\
&= (-1)^{N^2-1}\,\det(\mathcal{D}^{-1})\det(\tilde{\Delta}) \\
&= (-1)^{N^2-1}\,\det(\mathcal{D}^{-1})\det(\Delta^{-1})(\det(\partial_i D_i))^2,
\end{aligned} \tag{C.9}$$

where we have used the equation (B.3) and $\tilde{\Delta} = \partial\cdot D\Delta^{-1}\partial\cdot D$. Hence, we obtain that

$$\det\left(\begin{pmatrix} 0 & \mathbb{I}_{N^2-1} \\ \mathbb{I}_{N^2-1} & 0 \end{pmatrix} N'\right) = (-1)^{N^2-1}\,\det(\mathcal{D}^{-1})\det(\Delta^{-1})(\det(\partial_i D_i))^2. \tag{C.10}$$

On the other hand, it is easy to see that

$$\det\big(\delta_{kl} - \partial_k\Delta^{-1}\partial_l\big) = \det\begin{pmatrix} 1-\Delta^{-1}(\partial_1)^2 & -\Delta^{-1}\partial_1\partial_2 \\ -\Delta^{-1}\partial_2\partial_1 & 1-\Delta^{-1}(\partial_2)^2 \end{pmatrix} = \det(\Delta^{-1}(\partial_3)^2), \tag{C.11}$$

so that we have

$$\begin{aligned}
\det\left(\begin{pmatrix} 0 & \mathbb{I}_{N^2-1} \\ \mathbb{I}_{N^2-1} & 0 \end{pmatrix} N'\right) &= (-1)^{N^2-1}\,\det(N') \overset{(B.6)}{=} (-1)^{N^2-1}\,\det\Big(\big(\delta_{kl} - \partial_k\Delta^{-1}\partial_l\big)N\Big) \\
&= (-1)^{N^2-1}\,\det(\Delta^{-1}(\partial_3)^2)\det(N)
\end{aligned} \tag{C.12}$$

From Eqs.(C.10) and (C.12), we obtain that

$$\det(N) = \frac{\det(\mathcal{D}^{-1})(\det(\partial_i D_i))^2}{\det((\partial_3)^2)} \tag{C.13}$$

We have proved the equation (5.22) in the text.

References

[1] K. G. Wilson, *Phys. Rev.* **D10** (1974) 2445.

[2] D. J. Gross, R. D. Pisarski and L. G. Yaffe, *Rev. Mod. Phys.* **53**, 43 (1981).

[3] L. D. McLerran and B. Svetitsky, *Phys. Rev.* **D24** (1981) 450.

[4] M. Sakamoto and K. Takenaga, *Phys. Rev.* **D76** 085016 (2007); *Phys. Rev.* **D89** 105032 (2014).

[5] N. Weiss, *Phys. Rev.* **D24** (1981) 475; *Phys. Rev.* **D25** (1982) 2667.

[6] C. P. Korthals and M. Laine, *Phys. Lett.* **B511**, 269 (2001).

[7] K. Farakos and P. Pasipoularides, *Nucl. Phys.* **B705** (2005) 92 .

[8] Y. Hosotani, *Phys. Lett.* **B126** (1983) 309; *Ann. Phys. (N.Y.)* **190**, 233 (1989).

[9] V. Gribov, *Nucl. Phys.* **B139** (1978) 1.

[10] For a review see, G. Leibbrandt, *Rev. Mod. Phys.* **59**, 1067 (1987).

[11] W. Kummer, *Acta Phys. Austriaca* **14**, 149 (1961); R. L. Arnowitt and S. I. Fickier, *Phys. Rev.* **127**, 1821 (1962).

[12] A. M. Polyakov, *Phys. Lett.* **B72**, 447 (1978); J. Goldstone and R. Jackiw, *Phys. Lett.* **B74**, 81 (1978); V. Baluni and B. Grossman, *Phys. Lett.* **B78**, 226 (1978).

[13] L. Dolan and R. Jackiw, *Phys. Rev.* **D9** (1974) 3320; S. Weinberg, *Phys. Rev.* **D9** (1974) 3357.

[14] H. Reinhardt and J. Heffner, *Phys. Rev.* **D88** 045024 (2013).